



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The molecular basis of MeCP2 function in the brain

Citation for published version:

Tillotson, R & Bird, A 2020, 'The molecular basis of MeCP2 function in the brain', *Journal of Molecular Biology*, vol. 432, no. 6. <https://doi.org/10.1016/j.jmb.2019.10.004>

Digital Object Identifier (DOI):

[10.1016/j.jmb.2019.10.004](https://doi.org/10.1016/j.jmb.2019.10.004)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of Molecular Biology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





The Molecular Basis of MeCP2 Function in the Brain

Rebekah Tillotson^{1,2} and Adrian Bird³

1 - Genetics and Genome Biology Program, The Hospital for Sick Children, The Peter Gilgan Centre for Research and Learning, Toronto, ON M5G 0A4, Canada

2 - Medical Research Council (MRC) Molecular Haematology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Headington, Oxford, OX3 9DS, UK

3 - Wellcome Centre for Cell Biology, University of Edinburgh, The Michael Swann Building, King's Buildings, Max Born Crescent, Edinburgh, EH9 3BF, UK

Correspondence to Adrian Bird: a.bird@ed.ac.uk

<https://doi.org/10.1016/j.jmb.2019.10.004>

Edited by Tuncay Baubec

Abstract

MeCP2 is a reader of the DNA methylome that occupies a large proportion of the genome due to its high abundance and the frequency of its target sites. It has been the subject of extensive study because of its link with 'MECP2-related disorders', of which Rett syndrome is the most prevalent. This review integrates evidence from patient mutation data with results of experimental studies using mouse models, cell lines and *in vitro* systems to critically evaluate our understanding of MeCP2 protein function. Recent evidence challenges the idea that MeCP2 is a multifunctional hub that integrates diverse processes to underpin neuronal function, suggesting instead that its primary role is to recruit the NCoR1/2 co-repressor complex to methylated sites in the genome, leading to dampening of gene expression.

© 2019 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Numerous chromatin-associated proteins and noncoding RNAs work together to establish cell type-specific epigenetic states that regulate gene expression. These factors are vital for normal mammalian development, with deletion of individual members often resulting in lethality in mutant mice [1]. Additionally, epigenetic factors often have links to neurological disease, caused by mutations affecting dosage such as haploinsufficiency, locus duplications or hypomorphic alleles [2]. Here, we discuss the role of MeCP2, a reader of the DNA methylome, encoded by the X-linked *MECP2* gene. The gene has been implicated in several 'MECP2-related disorders' [3] prompting numerous studies of MeCP2 protein function. We discuss a spectrum of evidence that sheds light on the molecular mechanisms involved, including clinical genetic investigations of genotype-phenotype correlations and mouse models of the resulting human conditions.

Next-generation sequencing technologies have revealed where MeCP2 binds in the genome and how it interprets the DNA methylome to influence gene expression. Finally, structural studies have defined in molecular detail the interactions between MeCP2 and its key binding partners. Together, the findings provide a coherent picture of MeCP2 function as an essential reader of the DNA methylome in the brain that optimises neuronal transcription programmes.

MECP2-Related Disorders and Mouse Models

Loss of function mutations in the *MECP2* gene in hemizygous male patients cause neonatal encephalopathy, which is usually fatal before the age of 2 years [4]. The same mutations cause the severe neurological disorder Rett syndrome (RTT) in heterozygous females [5]. RTT occurs in 1 in

10–15,000 live female births [6], making it one of the most common causes of monogenic intellectual disability in females. The condition is characterised by 6–18 months of normal development before cognitive regression. Affected individuals lose learned speech and purposeful hand movements, develop stereotypies such as ‘hand-wringing’ and acquire motor deficiencies including gait ataxia [3]. Symptoms can also include microcephaly [7], respiratory problems [8] and seizures [9]. Disease progression plateaus and although patients show slow regression in later life, 70% reach the age of 50 [10]. Other *MECP2* mutations that may retain more protein function have been implicated in milder psychiatric disorders, such as autism and schizophrenia in both males and females [<http://mecp2.chw.edu.au/>]. The importance of correct MeCP2 dosage is evidenced by *MECP2* duplication syndrome, which doubles MeCP2 levels and mostly affects males as the locus containing *MECP2* is almost always copied within the X chromosome. This condition can run in families, with carrier females developing no or mild symptoms because of extreme (>90%) skewing of X chromosome inactivation, thereby silencing the duplicated locus [11]. Affected boys display some symptoms that overlap with RTT including intellectual disability, impaired language, gait abnormalities and seizures. Individuals with this condition suffer from recurrent infections because of immunological dysfunction, often leading to death at around 25 years of age [11,12]. Both RTT and *MECP2* duplication syndrome have been extensively modelled in mice, which display overt neurological defects as well as phenotypes that can be assessed by behavioural testing (Table 1) [13–17]. Such studies have established that MeCP2 is required for maintenance of neuronal function since deletion of the gene in adult mice causes RTT-like defects [18,19]. Excitingly, reactivation of *Mecp2* in symptomatic MeCP2-deficient mice leads to phenotypic reversal, indicating that development in the absence of MeCP2 causes little or no lasting damage [20,21]. Similarly, the behavioural phenotypes of mice overexpressing MeCP2 can be rescued by genetic deletion or antisense oligonucleotide-mediated downregulation of one copy [22]. These findings provide hope that both disorders will be curable.

MeCP2 is an Essential Reader of DNA Methylation in the Brain

MeCP2 was initially discovered over quarter of a century ago because of its ability to bind DNA in a methylation-specific manner [23]. The ~90 amino acid region responsible for binding was called the methyl-CpG binding domain (MBD) [24]. Sequence similarity searches using the MBD sequence identi-

fied 10 other proteins that contain this domain: MBD1–4, MBD5 (alternative names TAM1, KIAA1461), MBD6 (alternative names TAM2, KIAA1887), BAZ2A (alternative name TIP5), BAZ2B, SETDB1 (alternative names ESET, KMT1E) and SETDB2 (alternative name CLLD8) [25–32]. Of these, only MeCP2, MBD1, MBD2 and MBD4 have been shown to bind DNA *in vitro* in a methylation-specific manner [23,26,32,33]. Specificity for the same binding sites in DNA raises the possibility of competition between these proteins, though analysis of their temporal-spatial expression patterns reveals differences between cell types. MeCP2 is expressed in all tissues but reaches near-histone abundance in neurons ($\sim 16 \times 10^6$ molecules per nucleus) [34]. Its levels increase during embryonic and postnatal development, plateauing at 10 years in humans [35] and 5 weeks in mice [34]. MBD1 is expressed during neurogenesis but is then downregulated [36]. MBD2 and MBD4 are more widely expressed across somatic tissues, and MBD4 is the only family member detected in embryonic stem cells [36,37]. Whereas complete deletion of *Mecp2* in mice leads to severe neurological symptoms and death around 9 weeks of age [13,14], knocking out the other members results in minimal phenotypes [36,38–43]. Even though loss of MBD1 causes decreased neurogenesis, the animals have a normal lifespan and only mild behavioural defects [38,39]. These results suggest that MeCP2 is the MBD family member with the greatest role in interpreting the DNA methylome in the brain.

MeCP2 protein levels in human and mouse tissues correlate poorly with transcription of the *MECP2/Mecp2* gene [35]. This may be explained by the activity of several posttranscriptional regulatory mechanisms, including alternative splicing, use of different polyadenylation sites and posttranslational modification. The gene spans four exons, which are transcribed and spliced to form two isoforms, e1 and e2, where only e2 includes exon 2 (Fig. 1) [44,45]. Isoform e1 is the ancestral form, conserved across vertebrates, whereas isoform e2 is only present in mammals. The two isoforms are very similar, differing only at the extreme N-terminus, and most evidence indicates that they are functionally interchangeable [46]. Importantly, e2 mRNA is translated at a much lower efficiency because of the presence of an upstream ATG [44], so the great majority of MeCP2 protein in the brain is derived from e1. The gene contains four alternative polyadenylation sites, producing 3'UTRs ranging from 0.1 to 8.6 kb, although only the longest and shortest mRNAs are detected in neurons. The use of different polyadenylation sites determines whether they contain binding sites for proteins and miRNAs that regulate mRNA stability and translation [47–49]. MeCP2 protein function and stability is also thought to be affected by multiple posttranslational modifications,

Table 1. The RTT-like phenotypic signature in *Mecp2*-mutant hemizygous male mice.

Mutation (MGI)	Mutation type	Protein level	Median survival	Overt symptoms						Anxiety	Motor function
				Act.	Gait	HLC	Tr.	Br.	GC		
Null [13,95] (<i>Mecp2</i> ^{tm1.1Bird})	Δex3-4	None	9 wks	↓	+	+	+	+	+	↓/↑*	↓
Null [14,86,179] (<i>Mecp2</i> ^{tm1.1Jae})	Δex3	V. low (internal deletion)	6–12 wks	↓	+	+	+	+	+	↓	↓
Null [180] (<i>Mecp2</i> ^{tm1Ppl})	Δex3+part of ex4	None	8 wks	↓	+	+	+	+	+	↓	↓
<i>Mecp2</i> fl/y; Nestin-Cre [13] (<i>Mecp2</i> ^{tm1Bird} ; Tg(Nes-cre)1Kln)	CNS KO (Δex3-4)	65% recombination in brain	6–12 wks	↓	+	+	ND	ND	+	ND	ND
<i>Mecp2</i> fl/y; Nestin-Cre [14] (<i>Mecp2</i> ^{tm1Jae} ; Tg(Nes-cre)1Atp)	CNS KO (Δex3)	> 90 % recombination in brain	6–12 wks	↓	ND	ND	+	+	+	ND	ND
<i>Mecp2</i> STOP/y; Nestin-Cre [56] (<i>Mecp2</i> ^{tm2Bird} ; Tg(Nes-cre)1Kln)	Peripheral KO (ΔSTOP in CNS)	<0.1× in periphery	>1 yr	(↓)	–	–	–	–	–	ND	↓
R106W-Tavi [87] (<i>Mecp2</i> ^{tm4.1Joez})	RTT: MBD missense	~0.1× (of WT-TAVI)	10 wks	Present, cumulative score						ND	ND
R111G-EGFP(tg on null background) [95] (Tg(MECP2*R111G/EGFP)1Hzo)	RTT: MBD missense	~1× (tg)	11 wks	ND	ND	ND	ND	ND	ND	ND	ND
Y120D [88] (<i>Mecp2</i> ^{tm1NInd})	RTT: MBD missense	~0.5×	14 wks	+	+	+	+	ND	+	NS*	↓
R133C-EGFP [15] (<i>Mecp2</i> ^{tm6.1Bird})	RTT: MBD missense	~0.6× (of WT-EGFP)	42 wks	↓	+	+	+	(+)	+	↓	(↓)
T158M-EGFP [15] (<i>Mecp2</i> ^{tm4.1Bird})	RTT: MBD missense	~0.3× (of WT-EGFP)	13 wks	↓	+	+	+	(+)	+	↓	↓
T158M-TAVI [87] (<i>Mecp2</i> ^{tm3.1Joez})	RTT: MBD missense	~0.25× (of WT-TAVI)	14 wks	Present, cumulative score						ND	ND
T158A [86] (<i>Mecp2</i> ^{tm1.1Joez})	RTT: MBD missense	~0.5×	16 wks	↓	+	+	?	+	?	↓	↓
R306C [107,181] (<i>Mecp2</i> ^{tm2.1Meg})	RTT: NID missense	~1×	~19 wks	Present, cumulative score						↑	↓
R306C-EGFP [15] (<i>Mecp2</i> ^{tm5.1Bird})	RTT: NID missense	~1× (of WT-EGFP)	30 wks	↓	+	+	+	+	+	↓	↓
R306C-EGFP(tg on null background) [95] (Tg(MECP2*R306C/EGFP)1Hzo)	RTT: NID missense	~1×	18 wks	↓	ND	ND	ND	ND	ND	↑*	↓
P225R [120] (<i>Mecp2</i> ^{tm8.1Bird})	RTT: ID missense	~0.22×	50 wks	↓	+	+	+	+	+	↓	↓
P322L [120] (<i>Mecp2</i> ^{tm9.1Bird})	RTT: CTD missense	~0.03×	9 wks	↓	+	+	+	+	+	↓	↓
R168X [182–184] (<i>Mecp2</i> ^{tm1.1Jtc})	RTT: ID nonsense	V. low	12 wks	↓	+	+	+	+	+	↓	↓
R255X [185] (<i>Mecp2</i> ^{tm1.1Irsf})	RTT: ID nonsense	Undetectable	9 wks	ND	ND	ND	ND	+	ND	↓	↓
R270X-EGFP(tg on null background) [123] (Tg(MECP2*R270X/GFP)#Hzo)	RTT: ID nonsense	~1× (tg)	12 wks	↓	+	+	+	+	+	ND	ND
G273X-EGFP(tg on null background) [123] (Tg(MECP2*G273X/GFP)#Hzo)	RTT: ID nonsense	~1× (tg)	29 wks	↓	+	+	+	+	+	ND	ND
G273X-EGFP [129] (<i>Mecp2</i> ^{em5Bird})	RTT: ID nonsense	~1×	36 wks	↓	+	+	+	+	+	ND	ND
S385PfsX6 [120] (<i>Mecp2</i> ^{tm10.1Bird})	RTT: CTD Truncating (Patient mutation L386HfsX5)	~0.1× (mRNA) ~0.45×	20 wks	↓	+	+	+	+	+	ND	ND
Floxed (exons 3 + 4) [13,18,91,92] (<i>Mecp2</i> ^{tm1Bird})	Hypomorphic	~0.5×	>1 yr	–	+	(+)	–	+	–	NS	↓
S80A [51] (<i>Mecp2</i> ^{tm2.1Jae})	Mutation of phosphorylation site	~1×	Normal lifespan	ND	ND	ND	ND	ND	ND	ND	↓
T308A [52] (<i>Mecp2</i> ^{tm3Meg})	Mutation of phosphorylation site	~1×	Normal lifespan	ND	ND	+	ND	ND	ND	ND	↓
S421A/S424A [52] (<i>Mecp2</i> ^{tm3.1Jae})	Mutation of phosphorylation sites	~1×	Normal lifespan	ND	ND	ND	ND	ND	ND	ND	↑
ΔAT-hook1 [122] (<i>Mecp2</i> ^{em1Smoc})	Deleted AT hook 1	~1–1.8×	>48 wks	↓	ND	–	–	ND	ND	↑	↓
G273X[ΔNLS]-EGFP [129] (<i>Mecp2</i> ^{em6Bird})	RTT ID nonsense + mutated NLS	~1×	41 wks (NS vs G273X-EGFP)	↓	+	+	+	+	+	ND	ND

ΔN-EGFP [130] (Mecp2 ^{em1Bird})	Testing dispensability	~1x	—	—	—	—	NS
ΔNC-EGFP [130] (Mecp2 ^{em7.1Bird})	Testing dispensability	~1x	—	—	—	—	NS
ΔNIC-EGFP [130] (Mecp2 ^{em2Bird})	Testing dispensability	~0.5x	↓	+	—	(+)	↓
A140V [139,140] (Mecp2 ^{1m1Vnat})	Intellectual disability in males:	ND (mRNA ~1x)	ND	ND	ND	ND	ND
MeCP2(tg) [16,22] (Tg(MECP2)1Hzo)	MBD missense Overexpression	~2x (tg)	↓	ND	(FP)	ND	↑
R306C-EGFP(tg on WT background) [95] (Tg(MECP2*R306C)EGFP)1Hzo)	OE rescued by mutating NID	~1x + WT	NS	ND	ND	ND	NS
Tau_Mecp2 [17] (homozygous on WT background) (Mapt ^{1m1(Mecp2)Jae})	Overexpression	5–9 fold	ND	+	ND	+	ND
Tau_Mecp2 [94] (homozygous on WT background) (Mapt ^{1m1(Mecp2)Bird})	Overexpression	~3.8x	Death before weaning	N/A	N/A	N / N / N / A	N/A
Tau_Mecp2 [R306C] [94] (homozygous on WT background) (Mapt ^{1m3(Mecp2*R306C)Bird})	OE rescued by mutating NID	~2.6x + WT	—	—	—	A A A	NS

RTT models display six overt symptoms: Act. = reduced spontaneous activity; Gait = abnormal gait; HLC = hindlimb clasping (FP = fore paw clasping); Tr. = tremor; Br. = breathing irregularities; GC = deterioration of general condition. RTT models display decreased anxiety, measured by Elevated Plus Maze or Elevated Zero Maze. *Light/Dark test was used instead in these studies, showing increased or unaltered anxiety. RTT models displayed impaired motor function, measured by one or more of the following: accelerating rotarod, running wheel, parallel rod footslip and hanging wire. RTT-like symptoms lead to premature death: median survival is given where reported. For non-RTT mutants, survival is stated as greater than the time point (e.g. 1yr) to which they were followed and <50% animals died. Some studies simply state 'normal lifespan'.

ND = not determined; NS = not significantly different from WT controls; N/A = not applicable due to lethality; Bracketed results = weak phenotype. ? = symptom not mentioned (only cumulative scores given); MGI = Mouse Genome Informatics; MBD = methyl-CpG binding domain; NID = NCoR1/2 interaction domain; CTD = C-terminal domain; NLS = nuclear localisation signal; EGFP = enhanced green fluorescent protein; RTT = Rett syndrome; ID = intellectual disability; CNS = central nervous system; WT = wild-type; KO = knock out; OE = overexpression; fl = floxed; tg = transgene wks = weeks; yr = year. The null background used to characterise transgenic mice was Mecp2^{tm1.1Bird}.

including neuronal activity-dependent phosphorylation (Fig. 2) [50–54]. Despite the existence of these posttranscriptional mechanisms, it is notable that cells cannot normalise MeCP2 protein levels to cope with even a doubling in gene dosage, as is the case in *MECP2* duplication syndrome.

The requirement for MeCP2 in different tissues has been tested using *Cre-loxP* technology to delete the gene in different cell types. Deletion of *Mecp2* in the central nervous system (CNS) using *Nestin-Cre* phenocopies *Mecp2*-null mice, showing that CNS dysfunction is the major contributor to RTT-like pathogenicity (Table 1) [13,14]. Studies focusing on the role of MeCP2 in different brain regions or neuronal subtypes using corresponding Cre drivers demonstrate that MeCP2-deficiency adversely affects each region or cell type tested [55]. Conditional *Mecp2* gene activation has also been used to determine the origin of clinical features of RTT that affect peripheral tissues. Mice with normal levels of MeCP2 in the CNS but severely lacking MeCP2 in all peripheral tissues are broadly healthy and do not display RTT-like phenotypes, including gait abnormalities and irregular breathing, demonstrating that these traits are due to loss of MeCP2 in the brain (Table 1) [56]. Exercise fatigue and bone fragility are the only symptoms attributable to loss of MeCP2 expression in the periphery. These findings support the notion that the major role of MeCP2 is to regulate neurological function.

MeCP2 Binds mCG Dinucleotides and mCAC Trinucleotides

The ability of MeCP2 to bind DNA containing methylated mCG dinucleotides was first identified by Southwestern assay [23] and has since been confirmed using a variety of *in vitro* techniques, including electrophoretic mobility shift assay (EMSA), surface plasmon resonance (SPR), fluorescence polarisation, isothermal titration calorimetry (ITC) and ¹H/¹⁵N-heteronuclear single-quantum correlation (HSQC) nuclear magnetic resonance spectroscopy [24,57–60]. *In vivo* binding has been visualised using immunohistochemistry or by fusion to a visible tag, taking advantage of the high concentration of mCG sites in pericentromeric repeats in mouse cells [35,61,62], and binding dynamics were quantified using fluorescence recovery after photobleaching (FRAP) [63–65]. More recent studies have focused on determining where in the mammalian genome MeCP2 binds using chromatin immunoprecipitation followed by next-generation sequencing (ChIP-seq). This has proved to be challenging because of both the high abundance of MeCP2




Fig. 1. Structure of the *MECP2* gene and its transcripts and resulting protein isoforms. Schematic diagram of the human *MECP2* gene comprising four exons. Above: Isoform e1 mRNA is formed by splicing together exons 1, 3 and 4. The protein is translated from exon 1. Below: Isoform e2 mRNA formed by splicing together all four exons. The protein is translated from exon 2. Alternative polyadenylation sites could potentially be used for either isoform. The open reading frame is shown in grey except for the different e1 and e2 N-termini (yellow and orange, respectively), the MBD (blue) and the NID (pink). The UTRs are shown in white. The RTT-causing mutation in e1 (A2V) is annotated in red. MBD = methyl-CpG binding domain; NID = NCoR1/2 interaction domain; RTT = Rett syndrome.

protein and the large number of potential binding sites. In the bulk genome, CG dinucleotides occur on average every 100 bp and are highly methylated. In contrast, 'CpG Islands' are enriched in CG dinucleotides and tend to be unmethylated [66–68], but they account for only 1–2% of the genome. Consistent with MeCP2 occupying a large fraction of mCG sites, this analysis results in a relatively featureless ChIP-seq signal that follows methylation density, interrupted by dramatically reduced binding at unmethylated CpG islands [34,69,70]. In spite of technical limitations, peak-calling algorithms have been successfully employed for 'summit analysis' using ChIP-seq and bisulphite sequencing data from the mouse frontal cortex to produce aggregate plots showing an enrichment of peaks over mCG sites, but not at unmethylated CG dinucleotides [71]. To investigate

the *in vivo* DNA binding footprint of MeCP2 further, assay for transposase-accessible chromatin using sequencing (ATAC-seq) was used to determine protected genomic regions around mCG sites in cultured human neurons [72]. The ATAC-seq signal from wild-type samples was divided by the equivalent signal in *Mecp2* knock-out samples, revealing a striking DNA binding footprint corresponding to the region protected by MeCP2 over mCG [72]. This DNA methylation-dependent footprint, which is absent at nonmethylated CG sites, is 11 bp in size, consistent with *in vitro* DNaseI footprinting [24] and MNase protection [73].

High levels of mCH (where H is A, C or T) have recently been discovered in the brain [74], primarily at CA dinucleotides. The highest level of this non-CG methylation is found at the trinucleotide CAC (~12%




Fig. 2. Posttranslational modifications of MeCP2. Schematic diagram of mouse MeCP2 protein (e2 isoform) annotated with the activity-dependent phosphorylation sites (above) and other posttranslational modifications (below) found in brain tissue or neuronal/glial cell lines. All activity-dependent sites are phosphorylated upon neuronal activity except S80, which is phosphorylated under basal conditions and dephosphorylated upon neuronal activity. All residues are conserved between human and mouse, except T441 (A443 in humans). Activity-dependent phosphorylation: S80, S86, T148/149, S164, S229, S274, T308, S421 (human residue 423), S424 (human residue 426) [50–52,97,155–158]; Other phosphorylation: S13 [158,159], S68, S70 [157], Y120 [160], T160 [156], S166 [97,156], S178 [97], S216 [53], S399 (human residue 401) [51,158]; Ubiquitination: K12, K82, K119, K130, K135, K233, K249, K256, K271, K321 [158]; SUMOylation: K223 [161], K412 (human residue 414) [162]; Acetylation: K305/307, K321 [158], K447 (human residue 449) [163]; Methylation: R162 [164]; GlcNAcylation: T434 (human residue 436), T440/441 (human residue 442/A443) [165]. Poly(ADP-ribosyl)-lation sites have not been mapped [166]. MBD = methyl-CpG binding domain; NID = NCOR1/2 interaction domain.

of CAC sites) [70], as indicated by motif analysis of non-CG methylation in mouse and human brain tissue [75,76]. This is likely because of the relaxed enzymatic specificity of the *de novo* DNA methyltransferase, DNMT3A [77]. Although the percentage of methylation at individual CA dinucleotides is low, the higher abundance of CA compared with CG in the genome means that the total number of mCA sites approaches that of mCG sites [70,74]. Interestingly, non-CG methylation accumulates during neuronal maturation at the same time as the build-up of MeCP2 [34,35,74], raising the possibility that MeCP2 is able to bind methylation in a non-CG context. EMSA analysis demonstrated that MeCP2 could bind a probe containing mCH [76] and subsequent *in vitro* studies narrowed the sequence preference first to mCA [71] and then to mCAC (and to a lesser extent mCAT) trinucleotides [70]. The requirement for adenine in the second position suggested that MeCP2 could recognise the methyl group of thymine on the complementary strand. Indeed, removal of this group by replacement of thymine with uracil (mCAC/GTG to mCAC/GUG) abolished the interaction [70]. Binding to mCAC and mCAT but not mCAA or mCAG was confirmed in cultured cells co-transfected with MeCP2 and methylated oligonucleotides and ChIP-seq data from mouse hypothalamus indicated a strong peak of MeCP2 binding over mCAC (but not mCAT) sites *in vivo* [70]. Binding to mCAC dramatically increases the number of MeCP2 binding sites in the genome and alters their distribution. Comparison between neuronal subtypes found greater differences in the patterns of mCH than mCG, suggesting that mCAC is a major contributor to neuronal cell type-specific patterns of MeCP2 binding [78,79]. In mice, these patterns are established in the first few weeks of life

when DNMT3A binding and mCH deposition occurs over the transcribed regions of lowly expressed genes [80].

The oxidised form of methyl-cytosine, hydroxymethylcytosine (hmC), is the product of the first step of the active demethylation pathway mediated by the ten-eleven translocation (Tet) enzymes [81]. Like mCH, hmC accumulates to uniquely high levels in postmitotic neurons [74,82], prompting the search for readers. *In vitro* analyses found that MeCP2 could bind to probes containing hmCAC (and to a lesser extent hmCAT) [60,70] but could not bind the predominant form of hydroxymethyl-cytosine, hmCG dinucleotides [57,59,70,71]. Co-transfection of cultured cells with MeCP2 and hydroxymethylated oligonucleotides confirmed the ability of MeCP2 to bind hmCAC sites [70]. Unlike hmCG, hmCAC sites are very rare in the mammalian brain [74], suggesting that the ability to bind hmCAC may not be of biological relevance. Interestingly, accumulation of hmCG in neurons by oxidation of mCG sites means that MeCP2 can no longer bind at these loci. This, plus the appearance of novel mCAC binding sites, will profoundly redistribute the profile of MeCP2 binding in the neuronal genome during postnatal maturation.

The crystal structure of the MBD of MeCP2 bound to a DNA molecule containing a central mCG site showed that binding is mediated by direct interactions between the nucleotides making up the mCG site and two arginine ‘fingers’, Arg111 and Arg133 [83]. The Arg111 sidechain is constrained by hydrogen bonding to Asp121, but Arg133 is relatively unconstrained. Computational modelling based on this structure predicted that the MBD can bind to mCAC or hmCAC-containing DNA in a single orientation by altering the position of Arg133 alone

[70]. Recent structural studies have broadly confirmed this hypothesis for mCAC [84]. The minor structural change required to accommodate mCAC, together with the likelihood that the structure of MeCP2 outside the globular MBD is predicted to be largely disordered [85], suggests that MeCP2 does not interpret mCG and mCAC sites differently.

DNA Binding is Essential for MeCP2 Protein Function

A functional MBD is vital for the role of MeCP2 as a reader of the methylome. Its biological importance is indicated by the large number of RTT-causing missense mutations that map to this domain, almost all of which impair binding (Table 2). Further experimentation is needed to verify whether other, less well-characterised mutations also impact binding. For example, RTT-causing mutations affecting three prolines, P101, P127 and P152 were reported to bind methylated DNA *in vivo*, because of their localisation at pericentromeric foci in fixed cells [62]. However, a DNA binding defect for these mutants cannot be ruled out as many mutants with defective DNA binding dynamics (quantified by FRAP) nevertheless showed localisation by this assay [65]. In addition, there is evidence that many RTT-causing mutations in the MBD reduce protein stability as knock-in mouse models carrying MBD mutations have reduced protein levels (Table 1) [15,85–88]. Several other mutants, including P101S and P152R, have been found to have reduced stability *in vitro*, measured by free energy changes of protein unfolding or dispersion of HSQC spectra (Table 2) [89,90]. It is likely that both disruption of DNA binding and instability contribute to the disease phenotype in MBD mutants, as reduced expression of the wild-type protein to similar levels caused a much milder phenotype (Table 1) [91,92], whereas increasing expression of MBD mutants improved health and survival [93,94]. The finding that transgenic mice expressing MeCP2[R111G] at wild-type levels phenocopy *Mecp2*-null mice proves that loss of DNA binding alone is sufficient to cause Rett syndrome [95]. Further underlining the importance of DNA binding for the clinical phenotype, MeCP2[R133C] retains more DNA binding than other MBD mutants [15] and gives the mildest average clinical symptom score [96].

MeCP2 protein can be posttranslationally modified at multiple sites by phosphorylation, ubiquitination, SUMOylation, acetylation, methylation, GlcNAcylation and poly(ADP-ribosyl)-lation (Fig. 2). The discovery of activity-dependent phosphorylation sites located within or close to the MBD led to the proposal that DNA binding by MeCP2 could be controlled by neuronal activity by addition or removal of phosphate at these sites. Upon membrane

depolarisation, Ser86, Thr148/Ser149 and Ser164 gain phosphorylation and Ser80 loses phosphorylation [50,51]. Biochemical analysis of mutants that prevent or mimic phosphorylation (e.g. S80A and S164D) suggest that phosphorylation changes at these sites impair DNA binding [51,97]. Knock-in mice carrying the S80A mutation display very mild RTT-like symptoms (Table 1) [51], although it is also possible that replacement of the highly conserved MBD residue Ser80 with alanine directly compromises function. Computational modelling based on the crystal structure predicts that phosphorylation of Ser164 impacts DNA binding [97]. Two activity-dependent phosphorylation sites outside the MBD have also been proposed to impair DNA binding: Ser421 and Ser424 (human residues Ser423 and Ser426) [51,98], and knock-in mice with alanine at both sites are reported to recapitulate some of the features of mutants overexpressing MeCP2 (Table 1) [51,98], consistent with a gain of function effect. However, analysis of the wild-type protein by genome-wide ChIP-seq failed to identify any regions of the genome with reduced MeCP2 binding upon depolarisation [99]. Significantly, no RTT missense mutations have been found at activity-dependent phosphorylation sites. At present, therefore, the functional role of their phosphorylation remains elusive.

The Key Interaction Partner of MeCP2 is the NCoR1/2 Co-repressor Complex

A prevalent view of MeCP2 molecular activity is that it is a multifunctional hub, involved in several cellular processes via its interactions with over 40 putative binding partners (Fig. 3) [100,101]. These roles include transcriptional repression, transcriptional activation, alternative splicing, chromatin remodelling and miRNA processing. An alternative view, based on studies of its effects on transcription, is that the most important role of MeCP2 is to inhibit gene expression. DNA methylation-dependent repression was demonstrated in early studies using transfected reporter constructs that were enzymatically modified at CG sites using a bacterial methyltransferase [13,102,103]. Similar experiments also demonstrated repression mediated by mCH sites [76]. Discovery of its repressive activity led to the expectation that MeCP2-deficiency would cause upregulation of a discrete set of target genes, but the failure to unambiguously identify these in MeCP2-deficient mice initially stymied the field. Loss of MeCP2 instead led to small expression changes in both directions at a large number of genes, leading to the belief that MeCP2 might function both to repress and activate transcription [104,105]. To distinguish between direct and indirect activity, recent studies have used ChIP-seq and

Table 2. RTT-causing missense mutations in the MBD affect DNA binding and/or protein stability.

RTT missense mutation	mCG binding <i>in vitro</i>	mCG binding <i>in vivo</i>	Position in structure (PDB: 3C2I [83])	Stability	Mouse model
D97Y	ND	Reduced [62]	Interaction with R106	ND	Not made
L100V	Reduced [90]	Partial [62,65,142]	Loop	Destabilised [89,90]	Not made
L100R	ND	ND	Loop	ND	Not made
P101R	ND	Binding [62,142]	Loop	ND	Not made
P101S	Binding [60]	Binding [62,142]	Loop	Destabilised [60]	Not made
P101H	ND	Binding [62,142,186]	Loop	ND	Not made
P101L	ND	Binding [62,142]	Loop	ND	Not made
R106W	Reduced [60,90,187–189]	Reduced [62,63,65,87,135,137,142,190,191]	Stabilises Asx-ST motif	Destabilised [60,87]	RTT-like [87]
R106Q	Reduced [90]	Reduced [62,192]	Stabilises Asx-ST motif	Destabilised [192]	Not made
R106L	ND	ND	Stabilises Asx-ST motif	ND	Not made
L108H	ND	ND	Stabilises Asx-ST motif	ND	Not made
R111G	Reduced [84,90,189]	Reduced [62,65,95,137,142,186]	DNA interface	ND	RTT-like [95]
Y120D	Reduced [193]	Reduced [62,88]	Loop	Destabilised [88]	RTT-like [88]
L124F	ND	Reduced [62]	Hydrophobic core	ND	Not made
P127L	ND	Binding [62,142]	Loop	ND	Not made
A131D	ND	ND	Hydrophobic core	ND	Not made
R133C	Reduced [15,84,90,187–189,194]	Partial [15,65,135,137,142]	DNA interface	Destabilised [15,89,194]	RTT-like [15]
R133P	ND	ND	DNA interface	ND	Not made
R133L	ND	Reduced [62,142,186]	DNA interface	ND	Not made
S134C	Reduced [90]	Binding [62,142]	DNA interface	Destabilised [89,90]	Not made
S134F	Reduced [58]	ND	DNA interface (Backbone)	ND	Not made
S134P	ND	ND	DNA interface (Backbone)	ND	Not made
K135E	ND	Reduced [62]	DNA interface (Backbone)	ND	Not made
L138S	ND	ND	Alpha helix	ND	Not made
P152R	Reduced [90]	Binding [62,142,186]	Loop	Destabilised [89,90]	Not made
F155S	Reduced [90,187,188]	Reduced [62,186,190]	Loop	Destabilised [194]	Not made
D156E	Reduced [90]	Reduced [62]	Asx-ST motif	Destabilised [89,90]	Not made
D156A	ND	ND	Asx-ST motif	ND	Not made
F157L	ND	ND	Asx-ST motif	ND	Not made
F157I	ND	ND	Asx-ST motif	ND	Not made
T158M	Reduced [83,90,187,188]	Reduced [15,62,65,87,93,142,190]	Asx-ST motif	Destabilised [15,60,87,93,194]	RTT-like [15,87,93]
T158A	Reduced [83,90]	Reduced [62,86,142]	Asx-ST motif	Destabilised [86]	RTT-like [86]
G161V	ND	ND	Asx-ST motif	ND	Not made

MBD = methyl-CpG binding domain; RTT = Rett syndrome; PDB = Protein Data Bank; ND = not determined.




Fig. 3. MeCP2 has been reported to interact with over 40 binding partners across its length. Schematic diagram showing the interaction sites of MeCP2 binding partners characterised by their proposed function when complexed with MeCP2. The minimal domain required for transcriptional repression of a reporter gene (the TRD 205–310 [102]) is indicated by the shaded red box. RTT-causing missense mutations (red) and neutral variants found in males on the ExAC database (black) are shown above. The hotspot in the C-terminus where RTT-causing truncations occur is indicated by a red arrow. Regions deleted in mice expressing truncated MeCP2 [130] (N = N-terminus, I = Intervening region and C = C-terminus) are shown by grey dotted lines. Interaction sites for: methylated DNA 78–162 [24]; AT-rich DNA 183–195 (hook 1) [121] and 257–272 (hook 2) [121,123]; DNA ('basic patch') 274–340 [95]; KPNA3 249–268 [128]; KPNA4 249–270 [128]; TBL1/TBLR1 (NCoR1/2 complex) 298–309 [119]; SIN3A 108–206 and 207–308 [103,107]; c-Ski 163–309 [106]; PU.1 1–162 and 163–309 [108]; YY1 202–255 [110]; TFIIB 204–310 [117]; CBF1 [109] (unmapped); PRMT6 [109] (unmapped); SP3 [109] (unmapped); SOX2 [111] (unmapped); BRAHMA [109] (unmapped); G9a [114] (unmapped); HLCS [115] (unmapped); ATRX [137] 108–169; HP1α/β (unmapped), requires phospho-S229 [158]; HP1γ 1–55 [131], requires phospho-S229 [158]; SMC1/3 [167] (unmapped, requires phospho-S229 [158]); Lamin B 163–201 [168]; DNMT1 77–161, 162–206 and 207–310 [169]; TET1 [170] (unmapped); CREB1 [105] (unmapped); MYCN [171] (unmapped); RNA 160–200 [171]; YB-1195–329 [133], requires phospho-S80 [158]; FBP11 311-CT [132]; HYPC 311-CT [132]; PRPF3 1–140 and 207–308 [172]; LEDGF 263–293 [173]; DHX9 263–269 [173]; TDP-43 [173] (unmapped); FUS [173] (unmapped); DGCR8 380-CT, requires phospho-S80 [134]; FOXG1, requires 1–9 (unique to isoform e2) [174]; HIPK2 [155] (unmapped); CDKL5 202-CT [175]; IKKα [176] (unmapped); p300 [163] (unmapped); SIRT1 [163] (unmapped); HMGB1 207-CT (motif 380–386) [134]; HTT [177] (unmapped); MBD2 163–309 [178]; MeCP2 (dimerisation) 163–309 [178]. MBD = methyl-CpG binding domain; NID = NCoR1/2 interaction domain; RTT = Rett syndrome; SNP = single nucleotide polymorphism.

RNA-seq data to relate the level of wild-type MeCP2 binding *in vivo* to changes in transcription at genes in *Mecp2-null* mouse brain [69,70]. Based on the effect of MeCP2 deficiency on expression, genes were divided into three categories: upregulated/unchanged/downregulated, and MeCP2-enrichment at genes in each category was compared. Data collected from both cerebral cortex and hypothalamus showed that MeCP2-enrichment was highest over upregulated genes, consistent with the notion that the protein normally inhibits expression directly. Importantly, a reciprocal effect was observed in mice over-expressing MeCP2, as downregulated genes now had the highest MeCP2-enrichment. Lower MeCP2-enrichment over genes that are downregulated when MeCP2 is absent and upregulated when MeCP2 is overexpressed indicates that MeCP2 does not directly activate their transcription but suggests instead that these transcriptional changes are an indirect consequence of altered MeCP2 levels. It is notable that gene expression changes correlate with the density of both mCG and mCAC, suggesting MeCP2 interprets both sites in the same way [70–72]. The model of MeCP2 as a global repressor fits with the high frequency of MeCP2 binding sites in the genome [70] matched by the large number of MeCP2 molecules per cell [34] and with the robust association between DNA methylation and transcriptional inhibition [67].

Of the large number of proposed MeCP2 interaction partners, 13 proteins have been suggested to mediate its role in transcriptional silencing. The first group of partners comprises the HDAC-containing NCoR1/2 [106,107] and SIN3A complexes [103] and three co-repressors that interact with these complexes (cSki [106], PU.1 [108] and CBF1 [109]). The second group is made up of the transcription factors, YY1 [110], SOX2 [111] or SP3 [112]. Third, evidence has been presented that MeCP2 recruits the histone methyltransferases PRMT6 [113], G9a [114] or HLCS [115] or the chromatin remodeller BRM [116]. Finally, MeCP2 was reported to inhibit transcription by binding TFIIB, preventing its incorporation into the preinitiation complex [117]. To date, the binding sites of only six of these (the NCoR1/2 complex, SIN3A, c-Ski, PU.1, YY1 and TFIIB) have been mapped to regions of MeCP2 (Fig. 3). To identify the domain required for MeCP2-mediated transcriptional repression, fragments were fused to the DNA binding domain of the yeast transcription factor GAL4, and residues 207–310 were found to be sufficient to repress a reporter gene regulated by GAL4 binding elements [102]. This region was named the transcriptional repression domain (TRD). As none of the mapped interaction sites lies wholly outside the TRD, knowledge of this domain does not intrinsically exclude responsibility of any of the 13 interaction partners for repression.

A common approach to predicting the biological importance of protein regions or residues relies on evolutionary conservation. MeCP2 is 95% identical in amino acid sequence between human and mouse. This level of conservation is greater than the average for all proteins (86.4% identity [118]) and therefore suggests that the entire amino acid sequence is functionally important. A strikingly different picture is seen when the missense mutations causing Rett syndrome are mapped onto the protein, as most are confined to two discrete domains [107]: the MBD and a second smaller cluster at the C-terminal end of the TRD (Fig. 3). The importance of these two domains for MeCP2 protein function is further supported by the pattern of variants identified in the human population, as recorded in the ExAC database. Mapping these variants showed changes to the protein sequence could be tolerated throughout the protein sequence but rarely in these two domains (Fig. 3). Interestingly, the C terminal cluster of RTT-causing mutations coincides with the region responsible for the interaction between the NCoR1/2 complex and MeCP2: residues 298–309, named the NCoR1/2 interaction domain (NID) [107,119]. RTT-causing mutations in this cluster (P302R, K304E, K305R and R306C) destroy this interaction and disrupt repressive activity [107]. The NID interacts with the WD40 domains of transducin-beta like 1 (TBL1) and TBL1-related (TBLR1), two paralogs that are core components of the NCoR1/2 complex. Co-crystallisation of the NID peptide with the WD40 domain of TBLR1 showed that all four MeCP2 residues mutated in RTT make extensive contacts with TBLR1 [119].

The most common RTT-causing missense mutation in the NID is R306C, responsible for 5% of all cases. Knock-in mice carrying this mutation display the same phenotypic signature as other RTT mouse models (Table 1) [13,15,107], confirming pathogenicity. In addition, transcriptional analysis shows the same patterns of dysregulation in R306C knock-in mice as the other models [71]. Taken together, this evidence shows that disruption of the interaction between MeCP2 and the NCoR1/2 complex is sufficient to cause both the neurological defects and transcriptional changes common to all RTT models. The presence of an activity-dependent phosphorylation site, T308, in the NID provides a potential mechanism by which NCoR1/2 complex recruitment by MeCP2 could be regulated by neuronal activity [52]. The importance of this site was demonstrated *in vivo* by the production of T308A knock-in mice. Cortical neurons derived from T308A mice had reduced induction of activity-dependent genes upon, consistent with retaining the co-repressor complex. These mice displayed very mild RTT-like symptoms including impaired motor function, rather than the gain of function phenotype that might be predicted (Table 1) [52].

Testing the Bridge Hypothesis: MeCP2 Recruits the NCoR1/2 Co-repressor Complex to Methylated DNA

The weight of evidence discussed so far supports a model where MeCP2 recruits the NCoR1/2 co-repressor complex to methylated sites on chromatin, requiring two functional domains: the MBD and the NID. In other words, MeCP2 acts as a bridge between DNA and the co-repressor. Although most RTT-causing missense mutations affect these two domains, three lie elsewhere in the protein: A2V (e1 isoform only, Fig. 1), P225R and P322L. There is also a major cluster of truncating mutations downstream of the NID, making up ~10% of cases of RTT (Fig. 3) [120]. Considering first the A2V mutation, evidence indicates that it impairs posttranslational processing of MeCP2. Normally, the N-terminal methionine and up to five downstream alanine residues are excised from newly synthesised MeCP2, followed by acetylation of the alanine now at the N-terminus. Replacement of the alanine in position 2 with valine reduces the efficiency of methionine aminopeptidase (the enzyme responsible for cleavage), leading to greatly reduced MeCP2 stability [54]. Mouse models of the two proline mutations, P225R and P332L, reproduce the RTT phenotype (Table 1). Both mutant proteins have reduced abundance in the brain and both are deficient in their ability to recruit TBL1 to pericentromeric heterochromatin [120]. Two of the most common patient mutations in the C terminal cluster were chosen for characterisation. The first, L386HfsX5, drastically reduced MeCP2 abundance leading to RTT-like symptoms in knock-in mice (equivalent mutation in mice is S385PfsX6). Pathogenicity of the second, P389X, was not initially recapitulated in the mouse model (equivalent mutation in mice is P384_S385insPL; P387X), but introduction of this patient mutation in the context of the human protein sequence demonstrated that it too leads to greatly reduced MeCP2 protein levels. Intriguingly, the C-terminal truncating mutations are the first RTT models found to have lower mRNA levels, suggesting transcript stability is affected when this region is mutated [120]. In summary, all RTT-causing mutations adversely affect the level of bridge function and do not therefore uncover unanticipated functional domains in MeCP2.

A more stringent test of the bridge hypothesis is to ask whether other characterised interaction sites in MeCP2 that lie outside the MBD and NID are dispensable for its function. The ExAC database shows many single nucleotide polymorphisms (SNPs) elsewhere in the protein indicating that these regions can be altered without pathogenic consequences [107]. Since mapped interaction sites tend to be large, however, it is not obvious that the

neutral SNPs could disrupt binding. Exceptions are the two AT hooks, which have short core consensus sequences. Indeed, the presence of destructive SNPs in healthy hemizygous males and heterozygous females suggests that the AT hooks are not required for MeCP2 protein function [121]. Deletion of AT hook 1 in hemizygous male knock-in mice has no effect on lifespan but leads to mild symptoms that may be caused by increased MeCP2 levels in some brain regions [122]. Taken together, these findings contradict the proposal that MeCP2 functions to compact chromatin via the MBD and both AT hooks [123]. The notion that MeCP2 compacts chromatin was first raised by *in vitro* studies [124], but recent analysis found that chromatin in neurons lacking MeCP2 is in fact more compact than wild-type [125], consistent with smaller nuclear size [126]. It is possible that absence of MeCP2 allows access for another compacting protein, as suggested by increased levels of histone H1 [34], although this hypothesis has been questioned [127]. Another characterised short functional domain is the bipartite nuclear localisation signal (NLS) [61] which binds to importins, KPNA3 and KPNA4 [128,129]. Absence of RTT mutations that inactivate the NLS suggests that this too is nonessential [129]. In fact, disruption of the NLS does not interfere with nuclear localisation and has no phenotypic consequences in mice (Table 1) [128,129]. Interestingly, the small size of MeCP2 allows it to enter the nucleus where it can be retained by binding to methylated DNA [129].

To critically assess the functional dispensability of large regions of MeCP2 protein outside the MBD and NID, a series of knock-in mice expressing truncated versions of MeCP2 were produced: Δ N, Δ NC and Δ NIC (where N = the region N-terminal to the MBD, C = the region C-terminal to the NID and I = the intervening region between the MBD and the NID; Fig. 3) [130]. Both Δ N and Δ NC were expressed at wild-type levels and hemizygous males expressing the mutant proteins were indistinguishable from wild-type littermates (Table 1). The dispensable N and C termini, which together make up half of the length of MeCP2, have been reported to contain interaction sites for multiple binding partners including the heterochromatin protein HP1 [131], the splicing factors FBP11, HYPC and YB-1 [132,133], and the miRNA processing factor DGCR8 [134] (Fig. 3). Δ NIC mice were also viable, surviving for at least a year, but displayed mild neurological defects (Table 1) that are at least partially attributable to the reduced protein levels in these mutants (~50% of wild-type) [130]. This illustrates that evolutionarily conserved regions of MeCP2 are not needed to prevent Rett syndrome, even though they are involved in molecular interactions. Taken together, the evidence strongly suggests that MeCP2 exerts a sole key function, which is to bridge methylated DNA and the TBL1/TBLR1 subunits of the NCoR1/2

complex. An important corollary of this model is that Rett syndrome is likely to be one disease with a single root cause.

The relative dispensability of two-thirds of MeCP2 leads us to ponder why these regions are so well-conserved throughout evolution. One theory is that the other regions are required for higher functions that were not detected in the phenotypic characterisation of the mice expressing the truncated alleles [130]. Looking at variants that fall in the grey area between no phenotypic consequence (male hemizygotes in the ExAC database) and complete loss of function (Rett syndrome in females and neonatal encephalopathy in males) may shed light on this issue. MeCP2 mutations have been implicated in intellectual disability in males, mild intellectual disability in females and in cognitive disorders such as autism and schizophrenia in both males and females. Missense mutations occur throughout the protein [<http://mecp2.chw.edu.au/>] and could hypothetically affect MeCP2 by causing subtle differences in bridge activity or protein levels or by impacting another role of MeCP2 that is required for higher cognitive function. There is evidence that regions outside the MBD contribute subtly to its interaction with DNA, as deletions or mutations in these regions affect binding and diffusion dynamics [63,64,121,135,136]. There is also evidence that these regions are involved in regulating protein stability, for example the intervening domain contains Ser216 whose phosphorylation is reported to modulate MeCP2 levels [53]. To date, no mutations outside the MBD or NID specifically disrupt the interaction with any other binding partner. This has, however, been suggested for the MBD mutation A140V, which initially did not appear to compromise DNA binding [137]. A140V is the most common MeCP2 mutation causing intellectual disability in boys and minor abnormalities in female carriers [138] and its milder phenotype compared with RTT-causing mutations is recapitulated in knock-in mice (Table 1) [139,140]. The mutation weakens the interaction between MeCP2 and ATRX [137], a chromatin remodeller also mutated in cases of severe intellectual disability [141]. However, recent reports of minor changes in DNA binding affinity of the A140V protein [90,142,143] indicate that more work is needed to determine the molecular causes of pathogenicity.

Excess NCoR1/2 Complex Recruitment is Required for Toxicity in *MECP2* duplication Syndrome

At the transcriptional level, two-fold overexpression of MeCP2 has the opposite effect compared with loss-of-function mutations. Genes enriched for MeCP2 binding are upregulated in *Mecp2*-null mice and downregulated in mice overexpressing MeCP2

[69,70]. This suggests that neurological toxicity caused by MeCP2 overexpression is because of excess recruitment of the NCoR1/2 co-repressor complex at these genomic regions. If so, mutating the NID in the additional copy/copies should alleviate the toxic effects. This possibility has been addressed using two independent mouse models of MeCP2 overexpression: (1) a PAC transgene containing the entire human locus [16] and (2) expression of mouse *Mecp2* isoform e2 from the *Mapt* locus fused to the first 31 amino acids of Tau [17]. The former, which express MeCP2 at two-fold wild-type levels, displayed neurological phenotypes in behavioural tests and developed late-onset overt symptoms with 30% mortality between 5 months and 1 year of age [16]. Introduction of R306C into the transgene abolished these phenotypic consequences [95]. This result was confirmed using the second model with a more extreme phenotype. Mice with 3.8-fold levels of wild-type MeCP2 failed to survive to weaning, but this was fully rescued by the introduction of the R306C mutation into Tau-MeCP2. Not only did the mice survive to adulthood, they remained healthy with no phenotype in behavioural tests (Table 1) [94]. Thus, a functional NID is essential to confer the toxicity caused by MeCP2 overexpression.

Breaking the Bridge by Mutating TBLR1 can Occasionally Cause Rett Syndrome

The Bridge hypothesis implies that mutations in the NCoR1/2 complex might also break the DNA-MeCP2-NCoR1/2 complex bridge, resulting in RTT. The core NCoR1/2 complex is 1–2 MDa in size and contains NCoR1, NCoR2, GPS2, HDAC3, TBL1 and TBLR1 subunits. Knockout mice lacking the genes encoding core subunits (except TBL1) have been produced and show that each is essential for embryonic development [144–148]. As the NCoR1/2 complex has other molecular functions including, for example, repression via nuclear receptors in the steroid/thyroid/retinoid superfamily, it is unsurprising that loss-of-function mutations in its components have a more severe phenotype than loss of MeCP2. To specifically abolish its role in MeCP2-mediated repression, its interaction surface would need to be mutated so that MeCP2 can no longer bind without affecting other interactions. The NCoR1/2 complex consists of a central TBL1/TBLR1 tetramer, with each TBL1/TBLR1 dimer bound to one molecule of NCoR1 or NCoR2. These in turn act as scaffolds for GPS2 and HDAC3 [149]. TBL1/TBLR1 tetramerisation involves their N-terminal domains, with their C-terminal WD40 domains located on the outside of this complex, suggesting the MeCP2 interaction surface can be specifically mutated. A key difference between mutating this interface within MeCP2 and within the NCoR1/2 complex is that both TBL1 and

Table 3. Missense mutations in the WD40 domain of TBLR1 cause intellectual disability.

Mutation	Clinical description	MeCP2 binding	NCoR1/2 complex binding	Reference
H213Q	Global DD, delayed speech and language, hypotonia, facial dysmorphism			DECIPHER 280701
L282P	ASD			[195,196]
A311P	Abnormality of the nervous system	Abolished		DECIPHER 322627
D369E	Severe ID, ASD, ADHD, facial dysmorphism		Yes, HDAC3	DECIPHER 260528
D370N	A: RTT-like criteria, including regression (loss of learned speech and replacement of purposeful hand movements with stereotypies) followed by stabilisation; gait abnormalities and breathing problems B: Not described C: Hypertelorism, seizures, feeding difficulties, impaired pain sensation			A [153]: B: DECIPHER 387266 C: DECIPHER 285549
D370Y	ID, seizures, facial dysmorphism, skeletal abnormalities, haemangioma			DECIPHER 261213
Y395C	Amold-Chiari malformation, ID, microcephaly, seizures			DECIPHER 273583
S417R	Moderate ID, microcephaly, facial dysmorphism			DECIPHER 303914
P444R	Global DD, ASD, facial dysmorphism	Abolished	Yes, HDAC3	DECIPHER 273334
Y446C	Pierpont syndrome: developmental delay, facial dysmorphism, abnormal fat distribution in distal limbs			[154]
G460D	ID, delayed speech and language, hyperactivity, facial dysmorphism, sleep disturbance			DECIPHER 255757

DD = Developmental delay; ID = intellectual disability; ASD = autism spectrum disorder; ADHD = attention deficit hyperactivity disorder; RTT = Rett syndrome.

TBLR1 paralogs can bind MeCP2, and disrupting mutations are unlikely to occur in both genes. Another consideration is that patients with Rett syndrome are mosaics due to X chromosome inactivation, with half their cells expressing the wild-type and half the mutant copy of the X-linked *MECP2* gene. The location of the gene encoding TBL1, *TBL1X*, on the X chromosome raised the possibility that mutations in this gene could have similar molecular consequences. However, the conditions for this scenario are not met as *TBL1X* is thought to be biallelically expressed [150] and TBL1 is not the major paralog in the brain—TBLR1 is in fact five times more abundant [151].

Analysis of disease-causing mutations in the *TBL* genes and their corresponding clinical outcome may illuminate the role of the NCoR1/2 complex in the brain. The DECIPHER database lists cases of developmental delay in both males and females caused by deletions and duplications in the genes encoding TBL1 and TBLR1, suggesting that TBL protein dosage impacts development. Furthermore, as is common for genes that escape X chromosome inactivation, dosage in males is maintained in humans and other primates by a Y-linked homolog, *TBL1Y*, and duplication of this gene also causes developmental delay [152, DECIPHER]. The DECIPHER database and case reports list numerous missense mutations in TBLR1 that cause developmental delay in heterozygotes. These are mostly located in the WD40 domain (Table 3), suggesting some may impact MeCP2 binding. Intriguingly, two patient mutations, D369E and P444R, were recently shown to abolish the bridge between TBLR1 and MeCP2, whilst retaining the ability of TBLR1 to pulldown HDAC3 (indicating that the complex remains intact) [119]. The challenge now is to compare the clinical characteristics of patients with TBLR1 WD40 domain mutations with RTT patients. Indeed, one patient with the D370N mutation was diagnosed with classical RTT, meeting all main and six out of 11 supporting criteria [153]. The Y446C mutation, which causes Pierpont syndrome, on the other hand is clinically distinct from RTT as patients lack the essential period of apparently normal development followed by regression [153,154]. It is therefore likely that loss of MeCP2 binding plays a role in the pathogenicity of some but not all of these mutations.

Concluding Remarks

There is now strong evidence that MeCP2 is an important reader of the DNA methylome in neurons. Loss-of-function mutations in *MECP2* that cause Rett syndrome in patients pinpoint the MBD (responsible for binding to methylated DNA) and the NID (responsible for binding to the NCoR1/2 co-repressor complex) as the key domains essential for

MeCP2 protein function. Analysis of the co-crystal structures of these interactions shows in atomic detail how the mutated residues contribute to binding to these macromolecules, and most mutations have been demonstrated experimentally to impair binding. The bridge model, which proposes that MeCP2 recruits the NCoR1/2 complex to methylated DNA, is supported by the observation that all RTT-causing mutations throughout the length of MeCP2 affect this function, and deletion of the regions outside these two domains has only mild phenotypic consequences. These findings significantly enhance our understanding of the underlying biology of *MECP2*-related disorders and link these molecular mechanisms to disease-causing mutations in *TBLR1*. After many years of research, recent findings now convey a coherent model for the role of MeCP2 in globally modulating gene expression, particularly in neurons. While there is evidently more to learn about the downstream metabolic consequences of MeCP2 dysfunction, the new knowledge promises to underpin efforts to devise therapies for RTT and other *MECP2*-related disorders.

Acknowledgements

The authors thank Jacky Guy, Jim Selfridge, Matthew Lyst and John Connelly for comments on the manuscript. R.T. is a Sir Henry Wellcome Postdoctoral Fellow (210913/Z/18/Z). A.B. is supported by a Wellcome Trust Centre grant (091580/Z/10/Z), a Wellcome Investigator Award (107930/Z/15/Z), a Rett Syndrome Research Trust Consortium grant, a European Research Council Advanced grant (EC 694295 *Gen-Epix*) and is a member of the Simons Initiative for the Developing Brain. The University of Edinburgh is a charitable body, registered in Scotland, with registration number SC005336.

Conflicts of Interest

None.

Received 3 September 2019;

Received in revised form 4 October 2019;

Accepted 5 October 2019

Available online 17 October 2019

Keywords

MeCP2;
Rett syndrome;
DNA methylation;
Transcriptional regulation;
Neurological function

References

- [1] T.F. Meehan, N. Conte, D.B. West, J.O. Jacobsen, J. Mason, J. Warren, C.K. Chen, I. Tudose, M. Relac, P. Matthews, N. Karp, L. Santos, T. Fiegel, N. Ring, H. Westerberg, S. Greenaway, D. Sneddon, H. Morgan, G.F. Codner, M.E. Stewart, J. Brown, N. Horner, M. Haendel, N. Washington, C.J. Mungall, C.L. Reynolds, J. Gallegos, V. Gailus-Durner, T. Sorg, G. Pavlovic, L.R. Bower, M. Moore, I. Morse, X. Gao, G.P. Tocchini-Valentini, Y. Obata, S.Y. Cho, J.K. Seong, J. Seavitt, A.L. Beaudet, M.E. Dickinson, Y. Herault, W. Wurst, M.H. De Angelis, K.C. Kent Lloyd, A.M. Flenniken, L.M.J. Nutter, S. Newbigging, C. McKelvie, M.J. Justice, S.A. Murray, K.L. Svenson, R.E. Braun, J.K. White, A. Bradley, P. Flicek, S. Wells, W.C. Skarnes, D.J. Adams, H. Parkinson, A.M. Mallon, S.D.M. Brown, D. Smedley, Disease model discovery from 3,328 gene knockouts by the international mouse phenotyping Consortium, *Nat. Genet.* 49 (2017) 1231–1238, <https://doi.org/10.1038/ng.3901>.
- [2] The Deciphering Developmental Disorders Study, Large-scale discovery of novel genetic causes of developmental disorders, *Nature* 519 (2015) 223–228, <https://doi.org/10.1038/nature14135>.
- [3] J.L. Neul, W.E. Kaufmann, D.G. Glaze, J. Christodoulou, A.J. Clarke, N. Bahi-Buisson, H. Leonard, M.E.S. Bailey, N.C. Schanen, M. Zappella, A. Renieri, P. Huppke, A.K. Percy, Rett syndrome: revised diagnostic criteria and nomenclature, *Ann. Neurol. Now.* 68 (2010) 944–950, <https://doi.org/10.1002/ana.22124>.
- [4] P. Kankirawatana, H. Leonard, C. Ellaway, J. Scurlock, A. Mansour, C.M. Makris, L.S. Dure IV, M. Friez, J. Lane, C. Kiraly-Borri, V. Fabian, M. Davis, J. Jackson, J. Christodoulou, W.E. Kaufmann, D. Ravine, A.K. Percy, Early progressive encephalopathy in boys and *MECP2* mutations, *Neurology* 67 (2006) 164–166, <https://doi.org/10.1212/01.wnl.0000223318.28938.45>.
- [5] R.E. Amir, I.B. Van den Veyver, M. Wan, C.Q. Tran, U. Francke, H.Y. Zoghbi, Rett syndrome is caused by mutations in X-linked *MECP2*, encoding methyl-CpG-binding protein 2, *Nat. Genet.* 23 (1999) 185–188, <https://doi.org/10.1038/13810>.
- [6] B. Hagberg, Rett's Syndrome: prevalence and impact on progressive severe mental retardation in girls, *Acta Paediatr. Scand.* 74 (1985) 405–408, <https://doi.org/10.1111/j.1651-2227.1985.tb10993.x>.
- [7] A.M. Kerr, J.B.P. Stephenson, Clinical topics Rett's syndrome in the west of Scotland, *Br. Med. J.* 291 (1985) 579–582, <https://doi.org/10.1136/bmj.291.6495.579>.
- [8] A.K. Percy, J.L. Neul, D.G. Glaze, K.J. Motil, S.A. Skinner, M. Omar Khwaja, H.-S. Lee, J.B. Lane, J.O. Barrish, F. Annese, L. McNair, J. Graham, K. Barnes, Rett syndrome diagnostic criteria: lessons from the natural history study, *Ann. Neurol.* 68 (2010) 951–955, <https://doi.org/10.1016/j.freeradbiomed.2008.10.025>. The.
- [9] B. Hagberg, J. Aicardi, K. Dias, O. Ramos, A progressive syndrome of autism, dementia, ataxia, and loss of purposeful hand use in girls: Rett's syndrome: report of 35 cases, *Ann. Neurol.* 14 (1983) 471–479, <https://doi.org/10.1097/00001577-198901020-00018>.
- [10] D.C. Tarquinio, W. Hou, J.L. Neul, W.E. Kaufmann, D.G. Glaze, K.J. Motil, S.A. Skinner, H.S. Lee,

- A.K. Percy, The changing face of survival in Rett syndrome and MECP2-related disorders, *Pediatr. Neurol.* 53 (2015) 402–411, <https://doi.org/10.1016/j.pediatrneurol.2015.06.003>.
- [11] H. Van Esch, MECP2 duplication syndrome, *Mol. Syndromol.* 2 (2011) 128–136, <https://doi.org/10.1159/000329580>.
 - [12] H. Van Esch, M. Bauters, J. Ignatius, M. Jansen, M. Raynaud, K. Hollanders, D. Lugtenberg, T. Bienvenu, L.R. Jensen, J. Ge, C. Moraine, P. Marynen, J. Fryns, G. Froyen, Duplication of the MECP2 region is a frequent cause of severe mental retardation and progressive neurological symptoms in males, *Am. J. Hum. Genet.* 77 (2005) 442–453, <https://doi.org/10.1086/444549>.
 - [13] J. Guy, B. Hendrich, M. Holmes, J.E. Martin, A. Bird, A mouse Mecp2-null mutation causes neurological symptoms that mimic Rett syndrome, *Nat. Genet.* 27 (2001) 322–326, <https://doi.org/10.1038/85899>.
 - [14] R.Z. Chen, S. Akbarian, M. Tudor, R. Jaenisch, Deficiency of methyl-CpG binding protein-2 in CNS neurons results in a Rett-like phenotype in mice, *Nat. Genet.* 27 (2001) 327–331, <https://doi.org/10.1038/85906>.
 - [15] K. Brown, J. Selfridge, S. Lagger, J. Connelly, D. De Sousa, S. Webb, J. Guy, C. Merusi, M. V Koerner, A. Bird, The molecular basis of variable phenotypic severity among common missense mutations causing Rett syndrome, *Hum. Mol. Genet.* 25 (2016) 558–570, <https://doi.org/10.1093/hmg/ddv496>.
 - [16] A.L. Collins, J.M. Levenson, A.P. Vilaythong, R. Richman, D.L. Armstrong, J.L. Noebels, J.D. Sweatt, H.Y. Zoghbi, Mild overexpression of MeCP2 causes a progressive neurological disorder in mice, *Hum. Mol. Genet.* 13 (2004) 2679–2689, <https://doi.org/10.1093/hmg/ddh282>.
 - [17] S. Luikenhuis, E. Giacometti, C.F. Beard, R. Jaenisch, Expression of MeCP2 in postmitotic neurons rescues Rett syndrome in mice, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 6033–6038, <https://doi.org/10.1073/pnas.0401626101>.
 - [18] H. Cheval, J. Guy, C. Merusi, D. De Sousa, J. Selfridge, A. Bird, Postnatal inactivation reveals enhanced requirement for MeCP2 at distinct age windows, *Hum. Mol. Genet.* 21 (2012) 3806–3814, <https://doi.org/10.1093/hmg/dds208>.
 - [19] C.M. McGraw, R.C. Samaco, H.Y. Zoghbi, Adult neural function requires MeCP2, *Science* 333 (2011) 186, <https://doi.org/10.1126/science.1206593>.
 - [20] J. Guy, J. Gan, J. Selfridge, S. Cobb, A. Bird, Reversal of neurological defects in a mouse model of Rett syndrome, *Science* 315 (2007) 1143–1147, <https://doi.org/10.1126/science.1138389>.
 - [21] L. Robinson, J. Guy, L. McKay, E. Brockett, R.C. Spike, J. Selfridge, D. De Sousa, C. Merusi, G. Riedel, A. Bird, S.R. Cobb, Morphological and functional reversal of phenotypes in a mouse model of Rett syndrome, *Brain* 135 (2012) 2699–2710, <https://doi.org/10.1093/brain/aws096>.
 - [22] Y. Sztainberg, H. Chen, J.W. Swann, S. Hao, B. Tang, Z. Wu, J. Tang, Y.-W. Wan, Z. Liu, F. Rigo, H.Y. Zoghbi, Reversal of phenotypes in MECP2 duplication mice using genetic rescue or antisense oligonucleotides, *Nature* 528 (2015) 123–126, <https://doi.org/10.1038/nature16159>.
 - [23] J.D. Lewis, R.R. Meehan, W.J. Henzel, I. Maurer-Fogy, P. Jeppesen, F. Klein, A. Bird, Purification, sequence, and cellular localization of a novel chromosomal protein that binds to methylated DNA, *Cell* 69 (1992) 905–914, [https://doi.org/10.1016/0092-8674\(92\)90610-o](https://doi.org/10.1016/0092-8674(92)90610-o).
 - [24] X. Nan, R.R. Meehan, A. Bird, Dissection of the methyl-CpG binding domain from the chromosomal protein MeCP2, *Nucleic Acids Res.* 21 (1993) 4886–4892, <https://doi.org/10.1093/nar/21.21.4886>.
 - [25] S.H. Cross, R.R. Meehan, X. Nan, A. Bird, A component of the transcriptional repressor MeCP1 shares a motif with DNA methyltransferase and HRX proteins, *Nat. Genet.* 16 (1997) 256–259, <https://doi.org/10.1038/nm0798-822>.
 - [26] B. Hendrich, A. Bird, Identification and characterization of a family of mammalian methyl-CpG binding proteins, *Mol. Cell. Biol.* 18 (1998) 6538–6547, <https://doi.org/10.1128/MCB.18.11.6538>.
 - [27] R. Strohner, A. Nemeth, P. Jansa, U. Hofmann-Rohrer, R. Santoro, G. Längst, I. Grummt, NoRC - a novel member of mammalian ISWI-containing chromatin remodeling machines, *EMBO J.* 20 (2001) 4892–4900, <https://doi.org/10.1093/emboj/20.17.4892>.
 - [28] H. Mabuchi, H. Fujii, G. Calin, H. Mabuchi, H. Fujii, G. Calin, H. Alder, M. Negrini, L. Rassenti, T.J. Kipps, F. Bullrich, C.M. Croce, Novel candidate genes for leukemogenesis at chromosome 13q14, a region commonly deleted in B-cell chronic lymphocytic leukemia deleted in B-cell chronic lymphocytic leukemia 1, *Cancer Res.* 61 (2001) 2870–2877.
 - [29] D.C. Schultz, K. Ayyanathan, D. Negorev, G.G. Maul, F.J. Rauscher III, SETDB1: a novel KAP-1-associated histone H3, lysine 9-specific methyltransferase that contributes to HP1-mediated silencing of euchromatic genes by KRAB zinc-finger proteins, *Genes Dev.* 16 (2002) 919–932, <https://doi.org/10.1101/gad.973302>.
 - [30] T.C. Roloff, H.H. Ropers, U.A. Nuber, Comparative study of methyl-CpG-binding domain proteins, *BMC Genomics* 4 (2003) 1, <https://doi.org/10.1186/1471-2164-4-1>.
 - [31] B. Hendrich, S. Tweedie, The methyl-CpG binding domain and the evolving role of DNA methylation in animals, *Trends Genet.* 19 (2003) 269–277, [https://doi.org/10.1016/S0168-9525\(03\)00080-5](https://doi.org/10.1016/S0168-9525(03)00080-5).
 - [32] Q. Du, P.-L. Luu, C. Stirzaker, S.J. Clark, Methyl-CpG-binding domain proteins: readers of the epigenome, *Epigenomics* 7 (2015) 1051–1073, <https://doi.org/10.2217/epi.15.39>.
 - [33] S. Laget, M. Joulie, F. Le Masson, N. Sasai, E. Christians, S. Pradhan, R.J. Roberts, P.A. Defossez, The human proteins MBD5 and MBD6 associate with heterochromatin but they do not bind methylated DNA, *PLoS One* 5 (2010), e11982, <https://doi.org/10.1371/journal.pone.0011982>.
 - [34] P.J. Skene, R.S. Illingworth, S. Webb, A. Kerr, K.D. James, D.J. Turner, R. Andrews, A.P. Bird, Neuronal MeCP2 is expressed at near histone-octamer levels and globally alters the chromatin state, *Mol. Cell* 37 (2010) 457–468, <https://doi.org/10.1016/j.molcel.2010.01.030.Neuronal>.
 - [35] M.D. Shahbazian, B. Antalffy, D.L. Armstrong, H.Y. Zoghbi, Insight into Rett syndrome: MeCP2 levels display tissue- and cell-specific differences and correlate with neuronal maturation, *Hum. Mol. Genet.* 11 (2002) 115–124, <https://doi.org/10.1093/hmg/11.2.115>.
 - [36] K.H. Wood, B.S. Johnson, S.A. Welsh, J.Y. Lee, Y. Cui, E. Krizman, E.S. Brodtkin, J.A. Blendy, M.B. Robinson, M.S. Bartolomei, Z. Zhou, Tagging methyl-CpG-binding domain proteins reveals different spatiotemporal expression and supports distinct functions, *Epigenomics* 8 (2016) 455–473, <https://doi.org/10.2217/epi-2015-0004>.
 - [37] B. Hendrich, U. Hardeland, H.H. Ng, J. Jiricny, A. Bird, The thymine glycosylase MBD4 can bind to the product of deamination at methylated CpG sites, *Nature* 401 (1999) 301–304, <https://doi.org/10.1038/35006691>.
 - [38] X. Zhao, T. Ueba, B.R. Christie, B. Barkho, M.J. McConnell, K. Nakashima, E.S. Lein, B.D. Eadie, A.R. Willhoite,

- A.R. Muotri, R.G. Summers, J. Chun, K.-F. Lee, F.H. Gage, Mice lacking methyl-CpG binding protein 1 have deficits in adult neurogenesis and hippocampal function, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 6777–6782, <https://doi.org/10.1073/pnas.1131928100>.
- [39] A.M. Allan, X. Liang, Y. Luo, C. Pak, X. Li, K.E. Szulwach, D. Chen, P. Jin, X. Zhao, The loss of methyl-CpG binding protein 1 leads to autism-like behavioral deficits, *Hum. Mol. Genet.* 17 (2008) 2047–2057, <https://doi.org/10.1093/hmg/ddn102>.
- [40] B. Hendrich, J. Guy, B. Ramsahoye, V. a Wilson, A. Bird, Closely related proteins MBD2 and MBD3 play distinctive but interacting roles in mouse development, *Genes Dev.* 15 (2001) 710–723, <https://doi.org/10.1101/gad.194101>.
- [41] O.J. Sansom, J. Berger, S.M. Bishop, B. Hendrich, A. Bird, A.R. Clarke, Deficiency of Mbd2 suppresses intestinal tumorigenesis, *Nat. Genet.* 34 (2003) 145–147, <https://doi.org/10.1038/ng1155>.
- [42] C.B. Millar, J. Guy, O.J. Sansom, J. Selfridge, E. MacDougall, B. Hendrich, P.D. Keightley, S.M. Bishop, A.R. Clarke, A. Bird, Enhanced CpG mutability and tumorigenesis in MBD4-deficient mice, *Science* 297 (2002) 403–405, <https://doi.org/10.1126/science.1073354>.
- [43] E. Wong, K. Yang, M. Kuraguchi, U. Werling, E. Avdievich, K. Fan, M. Fazzari, B. Jin, A.M.C. Brown, M. Lipkin, W. Edelmann, Mbd4 inactivation increases Cright-arrowT transition mutations and promotes gastrointestinal tumor formation, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 14937–14942, <https://doi.org/10.1073/pnas.232579299>.
- [44] S. Kriaucionis, A. Bird, The major form of MeCP2 has a novel N-terminus generated by alternative splicing, *Nucleic Acids Res.* 32 (2004) 1818–1823, <https://doi.org/10.1093/nar/gkh349>.
- [45] G.N. Mnatzakanian, H. Lohi, I. Munteanu, S.E. Alfred, T. Yamada, P.J.M. MacLeod, J.R. Jones, S.W. Scherer, N.C. Schanen, M.J. Friez, J.B. Vincent, B. a Minassian, A previously unidentified MECP2 open reading frame defines a new protein isoform relevant to Rett syndrome, *Nat. Genet.* 36 (2004) 339–341, <https://doi.org/10.1038/ng1327>.
- [46] B. Kerr, J. Soto C, M. Saez, A. Abrams, K. Walz, J.I. Young, Transgenic complementation of MeCP2 deficiency: phenotypic rescue of Mecp2-null mice by isoform-specific transgenes, *Eur. J. Hum. Genet.* 20 (2012) 69–76, <https://doi.org/10.1038/ejhg.2011.145>.
- [47] M.E. Klein, D.T. Lioy, L. Ma, S. Impey, G. Mandel, R.H. Goodman, Homeostatic regulation of MeCP2 expression by a CREB-induced microRNA, *Nat. Neurosci.* 10 (2007) 1513–1514, <https://doi.org/10.1038/nn2010>.
- [48] V.A. Gennarino, C.E. Alcott, C. Chen, A. Chaudhury, M.A. Gillentine, J.A. Rosenfeld, S. Parikh, J.W. Wheless, E.R. Roeder, D.D.G. Horovitz, E.K. Roney, J.L. Smith, S.W. Cheung, W. Li, J.R. Neilson, C.P. Schaaf, NUDT21-spanning CNVs lead to neuropsychiatric disease and altered MeCP2 abundance via alternative polyadenylation, *Elife* 4 (2015), e10782, <https://doi.org/10.7554/eLife.10782>.
- [49] D.C. Rodrigues, D.S. Kim, G. Yang, K. Zaslavsky, K.C.H. Ha, R.S.F. Mok, P.J. Ross, M. Zhao, A. Piekna, W. Wei, B.J. Blencowe, Q. Morris, J. Ellis, MECP2 is post-transcriptionally regulated during human neurodevelopment by combinatorial action of RNA-binding proteins and miRNAs, *Cell Rep.* 17 (2016) 720–734, <https://doi.org/10.1016/j.celrep.2016.09.049>.
- [50] Z. Zhou, E.J. Hong, S. Cohen, W. ning Zhao, H. yi H. Ho, L. Schmidt, W.G. Chen, Y. Lin, E. Savner, E.C. Griffith, L. Hu, J.A.J. Steen, C.J. Weitz, M.E. Greenberg, Brain-specific phosphorylation of MeCP2 regulates activity-dependent Bdnf transcription, dendritic growth, and spine maturation, *Neuron* 52 (2006) 255–269, <https://doi.org/10.1016/j.neuron.2006.09.037>.
- [51] J. Tao, K. Hu, Q. Chang, H. Wu, N.E. Sherman, K. Martinowich, R.J. Klose, C. Schanen, R. Jaenisch, W. Wang, Y.E. Sun, Phosphorylation of MeCP2 at Serine 80 regulates its chromatin association and neurological function, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 4882–4887, <https://doi.org/10.1073/pnas.0811648106>.
- [52] D.H. Ebert, H.W. Gabel, N.D. Robinson, N.R. Kastan, L.S. Hu, S. Cohen, A.J. Navarro, M.J. Lyst, R. Ekiert, A.P. Bird, M.E. Greenberg, Activity-dependent phosphorylation of MeCP2 threonine 308 regulates interaction with NCoR, *Nature* 499 (2013) 341–345, <https://doi.org/10.1038/nature12348>.
- [53] L.M. Lombardi, M. Zaghlula, Y. Sztainberg, S.A. Baker, T.J. Klisch, A.A. Tang, E.J. Huang, H.Y. Zoghbi, An RNA interference screen identifies druggable regulators of MeCP2 stability, *Sci. Transl. Med.* 9 (2017), eaaf7588, <https://doi.org/10.1126/scitranslmed.aaf7588>.
- [54] T.I. Sheikh, A. Martínez de Paz, S. Akhtar, J. Ausió, J.B. Vincent, MeCP2 isoform N-terminal modifications affect its degradation rate and are disrupted by the Ala2Val Rett mutation, *Hum. Mol. Genet.* 26 (2017) 4132–4141, <https://doi.org/10.1093/hmg/ddx300>.
- [55] J. Guy, H. Cheval, J. Selfridge, A. Bird, The role of MeCP2 in the brain, *Annu. Rev. Cell Dev. Biol.* 27 (2011) 631–652, <https://doi.org/10.1146/annurev-cellbio-092910-154121>.
- [56] P.D. Ross, J. Guy, J. Selfridge, B. Kamal, E. Tanner, T.H. Gilligwater, R.A. Jones, C.M. Loughrey, C.S. Mccarroll, M.E.S.B. Adrian, S. Cobb, Exclusive expression of MeCP2 in the nervous system distinguishes between brain and peripheral Rett syndrome-like phenotypes, *Hum. Mol. Genet.* 25 (2016) 4389–4404, <https://doi.org/10.1093/hmg/ddw269>.
- [57] V. Valinluck, H.-H. Tsai, D.K. Rogstad, A. Burdzy, A. Bird, L.C. Sowers, Oxidative damage to methyl-CpG sequences inhibits the binding of the methyl-CpG binding domain (MBD) of methyl-CpG binding protein 2 (MeCP2), *Nucleic Acids Res.* 32 (2004) 4100–4108, <https://doi.org/10.1093/nar/gkh739>.
- [58] M. Mellén, P. Ayata, S. Dewell, S. Kriaucionis, N. Heintz, MeCP2 binds to 5hmC enriched within active genes and accessible chromatin in the nervous system, *Cell* 151 (2012) 1417–14130, <https://doi.org/10.1016/j.cell.2012.11.022>.
- [59] H. Hashimoto, Y. Liu, A.K. Upadhyay, Y. Chang, S.B. Howerton, P.M. Vertino, X. Zhang, X. Cheng, Recognition and potential mechanisms for replication and erasure of cytosine hydroxymethylation, *Nucleic Acids Res.* 40 (2012) 4841–4849, <https://doi.org/10.1093/nar/gks155>.
- [60] M.J. Sperlazza, S.M. Bilinovich, L.M. Sinanan, F.R. Javier, D.C. Williams, Structural basis of MeCP2 distribution on non-CpG methylated and hydroxymethylated DNA, *J. Mol. Biol.* 429 (2017) 1581–1594, <https://doi.org/10.1016/j.jmb.2017.04.009>.
- [61] X. Nan, P. Tate, E. Li, A. Bird, DNA methylation specifies chromosomal localization of MeCP2, *Mol. Cell. Biol.* 16 (1996) 414–421, <https://doi.org/10.1128/mcb.16.1.414>.
- [62] S. Kudo, Y. Nomura, M. Segawa, N. Fujita, M. Nakao, C. Schanen, M. Tamura, Heterogeneity in residual function of MeCP2 carrying missense mutations in the methyl CpG

- binding domain, *J. Med. Genet.* 40 (2003) 487–493, <https://doi.org/10.1136/jmg.40.7.487>.
- [63] M. Marchi, A. Guarda, A. Bergo, N. Landsberger, C. Kilstrup-Nielsen, G.M. Ratto, M. Costa, Spatio-temporal dynamics and localization of MeCP2 and pathological mutants in living cells, *Epigenetics* 2 (2007) 187–197, <https://doi.org/10.4161/epi.2.3.5057>.
- [64] A. Kumar, S. Kamboj, B.M. Malone, S. Kudo, J.L. Twiss, K.J. Czymmek, J.M. LaSalle, N.C. Schanen, Analysis of protein domains and Rett syndrome mutations indicate that multiple regions influence chromatin-binding dynamics of the chromatin-associated protein MECP2 in vivo, *J. Cell Sci.* 121 (2008) 1128–1137, <https://doi.org/10.1242/jcs.016865>.
- [65] L. Schmiedeberg, P. Skene, A. Deaton, A. Bird, A temporal threshold for formaldehyde crosslinking and fixation, *PLoS One* 4 (2009), e4636, <https://doi.org/10.1371/journal.pone.0004636>.
- [66] A. Bird, M. Taggart, M. Frommer, O.J. Miller, D. Macleod, A fraction of the mouse genome that is derived from islands of nonmethylated, CpG-rich DNA, *Cell* 40 (1985) 91–99, [https://doi.org/10.1016/0092-8674\(85\)90312-5](https://doi.org/10.1016/0092-8674(85)90312-5).
- [67] A. Bird, DNA methylation patterns and epigenetic memory, *Genes Dev.* 16 (2002) 6–21, <https://doi.org/10.1101/gad.947102>.
- [68] A. Deaton, A. Bird, CpG islands and the regulation of transcription, *Genes Dev.* 25 (2011) 1010–1022, <https://doi.org/10.1101/gad.2037511.1010>.
- [69] B. Kinde, D.Y. Wu, M.E. Greenberg, H.W. Gabel, DNA methylation in the gene body influences MeCP2-mediated gene repression, *Proc. Natl. Acad. Sci. U. S. A.* 113 (2016) 15114–15119, <https://doi.org/10.1073/pnas.1618737114>.
- [70] S. Lagger, J.C. Connelly, G. Schweikert, S. Webb, J. Selfridge, B.H. Ramsahoye, M. Yu, C. He, G. Sanguinetti, L.C. Sowers, M.D. Walkinshaw, A. Bird, MeCP2 recognizes cytosine methylated tri-nucleotide and di-nucleotide sequences to tune transcription in the mammalian brain, *PLoS Genet.* 13 (2017), e1006793, <https://doi.org/10.1371/journal.pgen.1006793>.
- [71] H.W. Gabel, B. Kinde, H. Stroud, C.S. Gilbert, D.a. Harmin, N.R. Kastan, M. Hemberg, D.H. Ebert, M.E. Greenberg, Disruption of DNA-methylation-dependent long gene repression in Rett syndrome, *Nature* 522 (2015) 89–93, <https://doi.org/10.1038/nature14319>.
- [72] J. Cholewa-waclaw, R. Shah, S. Webb, K. Chhatbar, B. Ramsahoye, O. Pusch, Quantitative modelling predicts the impact of DNA methylation on RNA polymerase II traffic, *Proc. Natl. Acad. Sci. U. S. A.* 116 (2019) 14995–15000, <https://doi.org/10.1073/pnas.1903549116>.
- [73] T. Nikitina, R.P. Ghosh, R.a. Horowitz-Scherer, J.C. Hansen, S.a. Grigoryev, C.L. Woodcock, MeCP2-Chromatin interactions include the formation of chromosome-like structures and are altered in mutations causing Rett syndrome, *J. Biol. Chem.* 282 (2007) 28237–28245, <https://doi.org/10.1074/jbc.M704304200>.
- [74] R. Lister, E. a Mukamel, J.R. Nery, M. Urich, C. a Puddifoot, N.D. Johnson, J. Lucero, Y. Huang, A.J. Dwork, M.D. Schultz, M. Yu, J. Tonti-Filippini, H. Heyn, S. Hu, J.C. Wu, A. Rao, M. Esteller, C. He, F.G. Haghighi, T.J. Sejnowski, M.M. Behrens, J.R. Ecker, Global epigenomic reconfiguration during mammalian brain development, *Science* 341 (2013) 1237905, <https://doi.org/10.1126/science.1237905>.
- [75] K.E. Varley, J. Gertz, K.M. Bowling, S.L. Parker, T.E. Reddy, F. Pauli-behn, M.K. Cross, B. a Williams, J. a Stamatoyannopoulos, G.E. Crawford, D.M. Absher, B.J. Wold, R.M. Myers, Dynamic DNA methylation across diverse human cell lines and tissues Dynamic DNA methylation across diverse human cell lines and tissues, *Genome Res.* 23 (2013) 555–567, <https://doi.org/10.1101/gr.147942.112>.
- [76] J.U. Guo, Y. Su, J.H. Shin, J. Shin, H. Li, B. Xie, C. Zhong, S. Hu, T. Le, G. Fan, H. Zhu, Q. Chang, Y. Gao, G.-L. Ming, H. Song, Distribution, recognition and regulation of non-CpG methylation in the adult mammalian brain, *Nat. Neurosci.* 17 (2014) 215–222, <https://doi.org/10.1038/nn.3607>.
- [77] B.H. Ramsahoye, D. Biniszkiewicz, F. Lyko, V. Clark, a P. Bird, R. Jaenisch, Non-CpG methylation is prevalent in embryonic stem cells and may be mediated by DNA methyltransferase 3a, *Proc. Natl. Acad. Sci. U. S. A.* 97 (2000) 5237–5242, <https://doi.org/10.1073/pnas.97.10.5237>.
- [78] A. Mo, E.A. Mukamel, F.P. Davis, C. Luo, G.L. Henry, S. Picard, M.A. Urich, J.R. Nery, T.J. Sejnowski, R. Lister, S.R. Eddy, J.R. Ecker, J. Nathans, Epigenomic signatures of neuronal diversity in the mammalian brain, *Neuron* 86 (2015) 1369–1384, <https://doi.org/10.1016/j.neuron.2015.05.018>.
- [79] W. Renthal, L.D. Boxer, S. Hrvatin, E. Li, A. Silberfeld, M.A. Nagy, E.C. Griffith, T. Vierbuchen, M.E. Greenberg, Characterization of human mosaic Rett syndrome brain tissue by single-nucleus RNA sequencing, *Nat. Neurosci.* 21 (2018), <https://doi.org/10.1038/s41593-018-0270-6>.
- [80] H. Stroud, S.C. Su, S. Hrvatin, A.W. Greben, W. Renthal, L.D. Boxer, M.A. Nagy, D.R. Hochbaum, B. Kinde, H.W. Gabel, M.E. Greenberg, Early-life gene expression in neurons modulates lasting epigenetic states, *Cell* 171 (2017) 1151–1164, <https://doi.org/10.1016/j.cell.2017.09.047>, e16.
- [81] D. Globisch, M. Münzel, M. Müller, S. Michalakakis, M. Wagner, S. Koch, T. Brückl, M. Biel, T. Carell, Tissue distribution of 5-hydroxymethylcytosine and search for active demethylation intermediates, *PLoS One* 5 (2010), e15367, <https://doi.org/10.1371/journal.pone.0015367>.
- [82] S. Kriaucionis, N. Heintz, The nuclear DNA base 5-hydroxymethylcytosine is present in Purkinje neurons and the brain, *Science* 325 (2014) 929–930, <https://doi.org/10.1126/science.1169786>.
- [83] K.L. Ho, I.W. McNae, L. Schmiedeberg, R.J. Klose, A.P. Bird, M.D. Walkinshaw, MeCP2 binding to DNA depends upon hydration at methyl-CpG, *Mol. Cell* 29 (2008) 525–531, <https://doi.org/10.1016/j.molcel.2007.12.028>.
- [84] M. Lei, W. Tempel, K. Liu, J. Min, Plasticity at the DNA recognition site of the MeCP2 mCG-binding domain, *Biochim. Biophys. Acta - Gene Regul. Mech.* 1862 (2019) 194409, <https://doi.org/10.1016/j.bbargm.2019.194409>.
- [85] R.P. Ghosh, T. Nikitina, R.A. Horowitz-scherer, L.M. Gierasch, V.N. Uversky, K. Hite, J.C. Hansen, L. Christopher, Unique physical properties and interactions of the domains of methylated DNA binding protein 2 (MeCP2), *Biochemistry* 49 (2010) 4395–4410, <https://doi.org/10.1021/bi9019753>.
- [86] D. Goffin, M. Allen, L. Zhang, M. Amorim, I.-T.J. Wang, A.-R.S. Reyes, A. Mercado-Berton, C. Ong, S. Cohen, L. Hu, J. a Blendy, G.C. Carlson, S.J. Siegel, M.E. Greenberg, Z. Zhou, Rett syndrome mutation MeCP2 T158A disrupts DNA binding, protein stability and ERP responses, *Nat.*

- Neurosci. 15 (2012) 274–283, <https://doi.org/10.1038/nn.2997>.
- [87] B.S. Johnson, Y.T. Zhao, M. Fasolino, J.M. Lamonica, Y.J. Kim, G. Georgakilas, K.H. Wood, D. Bu, Y. Cui, D. Goffin, G. Vahedi, T.H. Kim, Z. Zhou, Biotin tagging of MeCP2 in mice reveals contextual insights into the Rett syndrome transcriptome, *Nat. Med.* 23 (2017) 1203–1214, <https://doi.org/10.1038/nm.4406>.
- [88] A. Gandaglia, E. Brivio, S. Carli, M. Palmieri, F. Bedogni, G. Stefanelli, A. Bergo, B. Leva, C. Cattaneo, L. Pizzamiglio, M. Cicerone, V. Bianchi, C. Kilstup-Nielsen, I. D'Annessa, D. Di Marino, P. D'Adamo, F. Antonucci, A. Frasca, N. Landsberger, A novel Mecp2Y120D knock-in model displays similar behavioral traits but distinct molecular features compared to the Mecp2-null mouse implying precision medicine for the treatment of Rett syndrome, *Mol. Neurobiol.* 56 (2018) 4838–4854, <https://doi.org/10.1007/s12035-018-1412-2>.
- [89] T. Kucukkal, Y. Yang, O. Uvarov, W. Cao, E. Alexov, Impact of Rett syndrome mutations on MeCP2 MBD stability, *Biochemistry* 54 (2015) 6357–6368, <https://doi.org/10.1021/acs.biochem.5b00790>.
- [90] Y. Yang, T.G. Kucukkal, J. Li, E. Alexov, W. Cao, Binding analysis of methyl-CpG binding domain of MeCP2 and Rett syndrome mutations, *ACS Chem. Biol.* 11 (2016) 2706–2715, <https://doi.org/10.1021/acscchembio.6b00450>.
- [91] B. Kerr, M. Alvarez-Saavedra, M. a Sáez, A. Saona, J.I. Young, Defective body-weight regulation, motor control and abnormal social interactions in Mecp2 hypomorphic mice, *Hum. Mol. Genet.* 17 (2008) 1707–1717, <https://doi.org/10.1093/hmg/ddn061>.
- [92] R.C. Samaco, J.D. Fryer, J. Ren, S. Fyffe, H.T. Chao, Y. Sun, J.J. Greer, H.Y. Zoghbi, J.L. Neul, A partial loss of function allele of Methyl-CpG-binding protein 2 predicts a human neurodevelopmental syndrome, *Hum. Mol. Genet.* 17 (2008) 1718–1727, <https://doi.org/10.1093/hmg/ddn062>.
- [93] J.M. Lamonica, D.Y. Kwon, D. Goffin, P. Fenik, B.S. Johnson, Y. Cui, H. Guo, S. Veasey, Z. Zhou, Elevating expression of MeCP2 T158M rescues DNA binding and Rett syndrome-like phenotypes, *J. Clin. Invest.* 127 (2017) 1889–1904, <https://doi.org/10.1172/JCI90967>.
- [94] M. V Koerner, L. Fitzpatrick, J. Selfridge, J. Guy, D. De Sousa, R. Tillotson, A. Kerr, Z. Sun, M.A. Lazar, M.J. Lyst, A. Bird, Toxicity of overexpressed MeCP2 is independent of HDAC3 activity, *Genes Dev.* 32 (2018) 1514–1524, <https://doi.org/10.1101/gad.320325.118>.
- [95] L.D. Heckman, M.H. Chahrour, H.Y. Zoghbi, Rett-causing mutations reveal two domains critical for MeCP2 function and for toxicity in MECP2 duplication syndrome mice, *Elife* (2014), eLife2014, <https://doi.org/10.7554/eLife.02676.001> e02676.
- [96] V.A. Cuddapah, R.B. Pillai, K.V. Shekar, J.B. Lane, K.J. Motil, S.A. Skinner, D.C. Tarquinio, D.G. Glaze, G. McGwin, W.E. Kaufmann, A.K. Percy, J.L. Neul, M.L. Olsen, Methyl-CpG-binding protein 2 (MECP2) mutation type is associated with disease severity in Rett syndrome, *J. Med. Genet.* 51 (2014) 152–158, <https://doi.org/10.1136/jmedgenet-2013-102113>.
- [97] G. Stefanelli, A. Gandaglia, M. Costa, M.S. Cheema, D. Di Marino, I. Barbiero, C. Kilstup-Nielsen, J. Ausió, N. Landsberger, Brain phosphorylation of MeCP2 at serine 164 is developmentally regulated and globally alters its chromatin association, *Sci. Rep.* 6 (2016) 28295, <https://doi.org/10.1038/srep28295>.
- [98] H. Li, X. Zhong, K.F. Chau, E.C. Williams, Q. Chang, Loss of activity-induced phosphorylation of MeCP2 enhances synaptogenesis, LTP and spatial memory, *Nat. Neurosci.* 14 (2011) 1001–1008, <https://doi.org/10.1038/nn.2866>.
- [99] S. Cohen, H.W. Gabel, M. Hemberg, A.N. Hutchinson, L.A. Sadacca, D.H. Ebert, D.A. Harmin, R.S. Greenberg, V.K. Verdine, Z. Zhou, W.C. Wetsel, A.E. West, M.E. Greenberg, Genome-wide activity-dependent MeCP2 phosphorylation regulates nervous system development and function, *Neuron* 72 (2011) 72–85, <https://doi.org/10.1016/j.neuron.2011.08.022>.
- [100] F. Della Ragione, M. Vacca, S. Fioriniello, G. Pepe, M.D. Esposito, MECP2, a multi-talented modulator of chromatin architecture, *Brief. Funct. Genom.* 15 (2016) 420–431, <https://doi.org/10.1093/bfpg/elw023>.
- [101] M.J. Lyst, A. Bird, Rett syndrome: a complex disorder with simple roots, *Nat. Rev. Genet.* 16 (2015) 261–274, <https://doi.org/10.1038/nrg3897>.
- [102] X. Nan, F.J. Campoy, A. Bird, MeCP2 is a transcriptional repressor with abundant binding sites in genomic chromatin, *Cell* 88 (1997) 471–481, [https://doi.org/10.1016/S0092-8674\(00\)81887-5](https://doi.org/10.1016/S0092-8674(00)81887-5).
- [103] X. Nan, H.H. Ng, C.A. Johnson, C.D. Laherty, B.M. Turner, R.N. Eisenman, A. Bird, Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex, *Nature* 393 (1998) 386–389, <https://doi.org/10.1038/30764>.
- [104] M. Tudor, S. Akbarian, R.Z. Chen, R. Jaenisch, Nonlinear partial differential equations and applications: transcriptional profiling of a mouse model for Rett syndrome reveals subtle transcriptional changes in the brain, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 15536–15541, <https://doi.org/10.1073/pnas.242566899>.
- [105] M. Chahrour, S.Y. Jung, C. Shaw, X. Zhou, S.T.C. Wong, J. Qin, H.Y. Zoghbi, MeCP2, a key contributor to neurological disease, activates and represses transcription, *Science* 320 (2008) 1224–1229, <https://doi.org/10.1126/science.1153252>.
- [106] K. Kokura, S.C. Kaul, R. Wadhwa, T. Nomura, M.M. Khan, T. Shinagawa, T. Yasukawa, C. Colmenares, S. Ishii, The Ski protein family is required for MeCP2-mediated transcriptional repression, *J. Biol. Chem.* 276 (2001) 34115–34121, <https://doi.org/10.1074/jbc.M105747200>.
- [107] M.J. Lyst, R. Ekiert, D.H. Ebert, C. Merusi, J. Nowak, J. Selfridge, J. Guy, N.R. Kastan, N.D. Robinson, F. de Lima Alves, J. Rappsilber, M.E. Greenberg, A. Bird, Rett syndrome mutations abolish the interaction of MeCP2 with the NCoR/SMRT co-repressor, *Nat. Neurosci.* 16 (2013) 898–902, <https://doi.org/10.1038/nn.3434>.
- [108] M. Suzuki, T. Yamada, F. Kihara-Negishi, T. Sakurai, T. Oikawa, Direct association between PU.1 and MeCP2 that recruits mSin3A-HDAC complex for PU.1-mediated transcriptional repression, *Oncogene* 22 (2003) 8688–8698, <https://doi.org/10.1038/sj.onc.1207182>.
- [109] J. Mann, F. Oakley, F. Akiboye, a Elsharkawy, a W. Thorne, D. a Mann, Regulation of myofibroblast transdifferentiation by DNA methylation and MeCP2: implications for wound healing and fibrogenesis, *Cell Death Differ.* 14 (2007) 275–285, <https://doi.org/10.1038/sj.cdd.4401979>.
- [110] G. Forlani, E. Giarda, U. Ala, F. Di Cunto, M. Salani, R. Tupler, C. Kilstup-Nielsen, N. Landsberger, The

- MeCP2/YY1 interaction regulates ANT1 expression at 4q35: novel hints for Rett syndrome pathogenesis, *Hum. Mol. Genet.* 19 (2010) 3114–3123, <https://doi.org/10.1093/hmg/ddq214>.
- [111] K.E. Szulwach, X. Li, R.D. Smrt, Y. Li, Y. Luo, L. Lin, N.J. Santistevan, W. Li, X. Zhao, P. Jin, Cross talk between microRNA and epigenetic regulation in adult neurogenesis, *J. Cell Biol.* 189 (2010) 127–141, <https://doi.org/10.1083/jcb.200908151>.
- [112] C.K. Hwang, C.S. Kim, D.K. Kim, P. Law, L. Wei, H.H. Loh, Up-regulation of the mu-opioid receptor gene is mediated through chromatin remodeling and transcriptional factors in differentiated neuronal cells, *Mol. Pharmacol.* 78 (2010) 58–68, <https://doi.org/10.1124/mol.110.064311>.
- [113] S. Dhawan, S. Georgia, S. ing Tschen, G. Fan, A. Bhushan, Pancreatic beta cell identity is maintained by DNA methylation-mediated repression of *Arx*, *Dev. Cell* 20 (2011) 419–429, <https://doi.org/10.1016/j.devcel.2011.03.012>.
- [114] S. Subbanna, N.N. Nagre, M. Shivakumar, N.S. Umapathy, D. Psychoyos, B.S. Basavarajappa, Ethanol induced acetylation of histone at G9a exon1 and G9a-mediated histone H3 dimethylation leads to neurodegeneration in neonatal mice, *Neuroscience* 258 (2014) 422–432, <https://doi.org/10.1016/j.neuroscience.2013.11.043>.
- [115] J. Xue, S.S.K. Wijeratne, J. Zemleni, Holocarboxylase synthetase synergizes with methyl CpG binding protein 2 and DNA methyltransferase 1 in the transcriptional repression of long-terminal repeats, *Epigenetics* 8 (2013) 504–511, <https://doi.org/10.4161/epi.24449>.
- [116] K.N. Hari Krishnan, M.Z. Chow, E.K. Baker, S. Pal, S. Bassal, D. Brasacchio, L. Wang, J.M. Craig, P.L. Jones, S. Sif, A. El-Osta, Brahma links the SWI/SNF chromatin-remodeling complex with MeCP2-dependent transcriptional silencing, *Nat. Genet.* 37 (2005) 254–264, <https://doi.org/10.1038/ng1516>.
- [117] N.K. Kaludov, A.P. Wolffe, MeCP2 driven transcriptional repression in vitro: selectivity for methylated DNA, action at a distance and contacts with the basal transcription machinery, *Nucleic Acids Res.* 28 (2000) 1921–1928, <https://doi.org/10.1093/nar/28.9.1921>.
- [118] W. Makalowski, Mark S. Boguski, Evolutionary parameters of the transcribed mammalian genome: an analysis of 2,820 orthologous rodent and human sequences, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 9407–9412, <https://doi.org/10.1073/pnas.95.16.9407>.
- [119] V. Kruusvee, M.J. Lyst, C. Taylor, Ž. Tarnauskaitė, A.P. Bird, A.G. Cook, Structure of the MeCP2–TBLR1 complex reveals a molecular basis for Rett syndrome and related disorders, *Proc. Natl. Acad. Sci. U. S. A.* 114 (2017) E3243–E3250, <https://doi.org/10.1073/pnas.1700731114>.
- [120] J. Guy, B. Alexander-Howden, L. Fitzpatrick, D. Desousa, M. V Koerner, J. Selfridge, A. Bird, A mutation-led search for novel functional domains in MeCP2, *Hum. Mol. Genet.* 27 (2018) 2531–2545, <https://doi.org/10.1093/hmg/ddy159>.
- [121] M.J. Lyst, J. Connelly, C. Merusi, A. Bird, Sequence-specific DNA binding by AT-hook motifs in MeCP2, *FEBS Lett.* 590 (2016) 2927–2933, <https://doi.org/10.1002/1873-3468.12328>.
- [122] M. Xu, P. Song, W. Huang, R. He, Y. He, X. Zhou, Y. Gu, Disruption of AT-hook 1 domain in MeCP2 protein caused behavioral abnormality in mice, *Biochim. Biophys. Acta (BBA) - Mol. Basis Dis.* 1864 (2018) 347–358, <https://doi.org/10.1016/j.bbadis.2017.10.022>.
- [123] S.A. Baker, L. Chen, A.D. Wilkins, P. Yu, O. Lichtarge, H.Y. Zoghbi, An AT-hook domain in MeCP2 determines the clinical course of Rett syndrome and related disorders, *Cell* 152 (2013) 984–996, <https://doi.org/10.1016/j.cell.2013.01.038>.
- [124] P.T. Georgel, R.A. Horowitz-Scherer, N. Adkins, C.L. Woodcock, P.A. Wade, J.C. Hansen, Chromatin compaction by human MeCP2. Assembly of novel secondary chromatin structures in the absence of DNA methylation, *J. Biol. Chem.* 278 (2003) 32181–32188, <https://doi.org/10.1074/jbc.M305308200>.
- [125] M.W. Linhoff, S.K. Garg, G. Mandel, A high-resolution imaging approach to investigate chromatin architecture in complex tissues, *Cell* 163 (2015) 246–255, <https://doi.org/10.1016/j.cell.2015.09.002>.
- [126] M. Yazdani, R. Deogracias, J. Guy, R.A. Poot, A. Bird, Y. Barde, Disease modeling using embryonic stem cells: MeCP2 regulates nuclear size and RNA synthesis in neurons, *Stem Cells* 30 (2012) 2128–2139, <https://doi.org/10.1002/june>.
- [127] A. Ito-Ishida, H.K. Yamalanchilli, Y. Shao, S.A. Baker, L.D. Heckman, L.A. Lavery, J. Kim, L.M. Lombardi, Y. Sun, Z. Liu, H.Y. Zoghbi, Genome-wide distribution of linker histone H1.0 is independent of MeCP2, *Nat. Neurosci.* 21 (2018) 794–798, <https://doi.org/10.1038/s41593-018-0155-8>.
- [128] S.A. Baker, L.M. Lombardi, H.Y. Zoghbi, Karyopherin alpha 3 and karyopherin alpha 4 mediate nuclear import of methyl-CpG binding protein 2, *J. Biol. Chem.* 290 (2015) 22485–22493, <https://doi.org/10.1074/jbc.M115.658104>.
- [129] M.J. Lyst, R. Ekiert, J. Guy, J. Selfridge, M.V. Koerner, C. Merusi, D. De Sousa, A. Bird, Affinity for DNA contributes to NLS independent nuclear localization of MeCP2, *Cell Rep.* 24 (2018) 2213–2220, <https://doi.org/10.1016/j.celrep.2018.07.099>.
- [130] R. Tillotson, J. Selfridge, M.V. Koerner, K.K.E. Gadalla, J. Guy, D. De Sousa, R.D. Hector, S.R. Cobb, A. Bird, Radically truncated MeCP2 reverses Rett syndrome-like neurological defects, *Nature* 550 (2017) 398–401, <https://doi.org/10.1038/nature24058>.
- [131] N. Agarwal, T. Hardt, A. Brero, D. Nowak, U. Rothbauer, A. Becker, H. Leonhardt, M.C. Cardoso, MeCP2 interacts with HP1 and modulates its heterochromatin association during myogenic differentiation, *Nucleic Acids Res.* 35 (2007) 5402–5408, <https://doi.org/10.1093/nar/gkm599>.
- [132] J.P. Buschdorf, W.H. Strätling, A WW domain binding region in methyl-CpG-binding protein MeCP2: impact on Rett syndrome, *J. Mol. Med.* 82 (2004) 135–143, <https://doi.org/10.1007/s00109-003-0497-9>.
- [133] J.I. Young, E.P. Hong, J.C. Castle, J. Crespo-Barreto, A.B. Bowman, M.F. Rose, D. Kang, R. Richman, J.M. Johnson, S. Berget, H.Y. Zoghbi, Regulation of RNA splicing by the methylation-dependent transcriptional repressor methyl-CpG binding protein 2, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 17551–17558, <https://doi.org/10.1073/pnas.0507856102>.
- [134] T.-L. Cheng, Z. Wang, Q. Liao, Y. Zhu, W.-H. Zhou, W. Xu, Z. Qiu, MeCP2 suppresses nuclear microRNA processing and dendritic growth by regulating the DGCR8/Drosha complex, *Dev. Cell* 28 (2014) 547–560, <https://doi.org/10.1016/j.devcel.2014.01.032>.
- [135] F.M. Piccolo, Z. Liu, P. Dong, C.-L. Hsu, E.I. Stoyanova, A. Rao, R. Tjian, N. Heintz, Mecp2 nuclear dynamics in live neurons results from low and high affinity chromatin

- interactions, *BioRxiv* (2019), <https://doi.org/10.2139/ssrn.3376659>.
- [136] T.I. Sheikh, R. Harripaul, M. Ayub, J.B. Vincent, MeCP2 AT-Hook1 mutations in patients with intellectual disability and/or schizophrenia disrupt DNA binding and chromatin compaction in vitro, *Hum. Mutat.* 39 (2018) 717–728, <https://doi.org/10.1002/humu.23409>.
- [137] X. Nan, J. Hou, A. Maclean, J. Nasir, M.J. Lafuente, X. Shu, S. Kriaucionis, A. Bird, Interaction between chromatin proteins MECP2 and ATRX is disrupted by mutations that cause inherited mental retardation, *Proc. Natl. Acad. Sci. U. S. A.* 104 (2007) 2709–2714, <https://doi.org/10.1073/pnas.0608056104>.
- [138] J.L. Neul, The relationship of Rett syndrome and MECP2 disorders to autism, *Transl. Res.* 14 (2012) 253–262.
- [139] G.M. Jentarra, S.L. Olfers, S.G. Rice, N. Srivastava, G.E. Homanics, M. Blue, S. Naidu, V. Narayanan, Abnormalities of cell packing density and dendritic complexity in the MeCP2 A140V mouse model of Rett syndrome/X-linked mental retardation, *BMC Neurosci.* 11 (2010), <https://doi.org/10.1186/1471-2202-11-19>.
- [140] L.Y. Ma, C. Wu, Y. Jin, M. Gao, G.H. Li, D. Turner, J.X. Shen, S.J. Zhang, V. Narayanan, G. Jentarra, J. Wu, Electrophysiological phenotypes of MeCP2 A140V mutant mouse model, *CNS Neurosci. Ther.* 20 (2014) 420–428, <https://doi.org/10.1111/cns.12229>.
- [141] R.J. Gibbons, D.J. Picketts, L. Villard, D.R. Higgs, Mutations in a putative global transcriptional regulator cause X-linked mental retardation with α -thalassemia (ATR-X syndrome), *Cell* 80 (1995) 837–845, [https://doi.org/10.1016/0092-8674\(95\)90287-2](https://doi.org/10.1016/0092-8674(95)90287-2).
- [142] N. Agarwal, A. Becker, K.L. Jost, S. Haase, B.K. Thakur, A. Brero, T. Hardt, S. Kudo, H. Leonhardt, M.C. Cardoso, MeCP2 Rett mutations affect large scale chromatin organization, *Hum. Mol. Genet.* 20 (2011) 4187–4195, <https://doi.org/10.1093/hmg/ddr346>.
- [143] J.Y. Chia, W.S. Tan, C.L. Ng, N. Hu, H.L. Foo, A/T run geometry of B-form DNA is independent of bound methyl-CpG binding domain, cytosine methylation and flanking sequence, *Sci. Rep.* 6 (2016) 31212, <https://doi.org/10.1038/srep31210>.
- [144] K. Jepsen, O. Hermanson, T.M. Onami, A.S. Gleiberman, V. Lunyak, R.J. Mcevilly, R. Kurokawa, V. Kumar, F. Liu, E. Seto, S.M. Hedrick, G. Mandel, C.K. Glass, D.W. Rose, M.G. Rosenfeld, S. Brook, N. York, Combinatorial roles of the nuclear receptor corepressor in transcription and development, *Cell* 102 (2000) 753–763, [https://doi.org/10.1016/S0092-8674\(00\)00064-7](https://doi.org/10.1016/S0092-8674(00)00064-7).
- [145] K. Jepsen, D. Solum, T. Zhou, R.J. McEvelly, H.J. Kim, C.K. Glass, O. Hermanson, M.G. Rosenfeld, SMRT-mediated repression of an H3K27 demethylase in progression from neural stem cell to neuron, *Nature* 450 (2007) 415–419, <https://doi.org/10.1038/nature06270>.
- [146] S. Bhaskara, B.J. Chyla, J.M. Amann, S.K. Knutson, D. Cortez, Z. Sun, S.W. Hiebert, Article deletion of histone deacetylase 3 reveals critical roles in S phase progression and DNA damage control, *Mol. Cell* 30 (2008) 61–72, <https://doi.org/10.1016/j.molcel.2008.02.030>.
- [147] C. Guo, Y. Li, C. Gow, M. Wong, J. Zha, C. Yan, H. Liu, Y. Wang, T.P. Burris, J. Zhang, The optimal corepressor function of nuclear receptor corepressor (NCoR) for peroxisome proliferator-activated receptor γ requires G protein pathway suppressor 2, *J. Biol. Chem.* 290 (2015) 3666–3679, <https://doi.org/10.1074/jbc.M114.598797>.
- [148] M.E. Dickinson, A.M. Flenniken, X. Ji, L. Teboul, M.D. Wong, J.K. White, T.F. Meehan, W.J. Weninger, H. Westerberg, H. Adissu, C.N. Baker, L. Bower, J.M. Brown, L. Brianna Caddle, F. Chiani, D. Clary, J. Cleak, M.J. Daly, J.M. Denegre, B. Doe, M.E. Dolan, S.M. Edie, H. Fuchs, V. Gailus-Durner, A. Galli, A. Gambadoro, J. Gallegos, S. Guo, N.R. Horner, C. wei Hsu, S.J. Johnson, S. Kalaga, L.C. Keith, L. Lanoue, T.N. Lawson, M. Lek, M. Mark, S. Marschall, J. Mason, M.L. McElwee, S. Newbigging, L.M.J. Nutter, K.A. Peterson, R. Ramirez-Solis, D.J. Rowland, E. Ryder, K.E. Samocha, J.R. Seavitt, M. Selloum, Z. Szoke-Kovacs, M. Tamura, A.G. Trainor, I. Tudose, S. Wakana, J. Warren, O. Wendling, D.B. West, L. Wong, A. Yoshiki, D.G. MacArthur, G.P. Tocchini-Valentini, X. Gao, P. Flicek, A. Bradley, W.C. Skarnes, M.J. Justice, H.E. Parkinson, M. Moore, S. Wells, R.E. Braun, K.L. Svenson, M. Hrabe de Angelis, Y. Herault, T. Mohun, A.M. Mallon, R. Mark Henkelman, S.D.M. Brown, D.J. Adams, K.C. Kent Lloyd, C. McKelrie, A.L. Beaudet, M. Bucan, S.A. Murray, M. McKay, B. Urban, C. Lund, E. Froeter, T. LaCasse, A. Mehalow, E. Gordon, L.R. Donahue, R. Taft, P. Kutney, S. Dion, L. Goodwin, S. Kales, R. Urban, K. Palmer, F. Pertuy, D. Bitz, B. Weber, P. Goetz-Reiner, H. Jacobs, E. Le Marchand, A. El Amri, L. El Fertak, H. Ennah, D. Ali-Hadji, A. Ayadi, M. Wattenhofer-Donze, S. Jacquot, P. André, M.C. Birling, G. Pavlovic, T. Sorg, I. Morse, F. Benso, M.E. Stewart, C. Copley, J. Harrison, S. Joynson, R. Guo, D. Qu, S. Spring, L. Yu, J. Ellegood, L. Morikawa, X. Shang, P. Feugas, A. Creighton, P.C. Penton, O. Danisment, N. Griggs, C.L. Tudor, A.L. Green, C. Icoresi Mazzeo, E. Siragher, C. Lillistone, E. Tuck, D. Gleeson, D. Sethi, T. Bayzatinova, J. Burvill, B. Habib, L. Weavers, R. Maswood, E. Miklejewska, M. Woods, E. Grau, S. Newman, C. Sinclair, E. Brown, S. Ayabe, M. Iwama, A. Murakami, High-throughput discovery of novel developmental phenotypes, *Nature* 537 (2016) 508–514, <https://doi.org/10.1038/nature19356>.
- [149] J. Oberoi, L. Fairall, P.J. Watson, J.-C. Yang, Z. Czimmerer, T. Kampmann, B.T. Goult, J.A. Greenwood, J.T. Gooch, B.C. Kallenberger, L. Nagy, D. Neuhaus, J.W.R. Schwabe, Structural basis for the assembly of the SMRT/NCoR core transcriptional repression machinery, *Nat. Struct. Mol. Biol.* 18 (2011) 177–184, <https://doi.org/10.1038/nsmb.1983>.
- [150] M.T. Bassi, R.S. Ramesar, B. Caciotti, I.M. Winship, A. De Grandi, M. Riboni, P.L. Townes, P. Beighton, A. Ballabio, G. Borsani, X-linked late-onset sensorineural deafness caused by a deletion involving OA1 and a novel gene containing WD-40 repeats, *Am. J. Hum. Genet.* 64 (1999) 1604–1616, <https://doi.org/10.1086/302408>.
- [151] C.G. Spruijt, F. Gnerlich, A.H. Smits, T. Pfaffeneder, P.W.T.C. Jansen, C. Bauer, M. Münzel, M. Wagner, M. Müller, F. Khan, H.C. Eberl, A. Mensinga, A.B. Brinkman, K. Lephikov, U. Müller, J. Walter, R. Boelens, H. Van Ingen, H. Leonhardt, T. Carell, M. Vermeulen, Dynamic readers for 5-(Hydroxy)methylcytosine and its oxidized derivatives, *Cell* 152 (2013) 1146–1159, <https://doi.org/10.1016/j.cell.2013.02.004>.
- [152] H.T. Yan, T. Shinka, K. Kinoshita, Y. Sato, M. Umeno, G. Chen, K. Tsuji, Y. Unemi, X.J. Yang, T. Iwamoto, Y. Nakahori, Molecular analysis of TBL1Y, a Y-linked homologue of TBL1X related with X-linked late-onset

- sensorineural deafness, *J. Hum. Genet.* 50 (2005) 175–181, <https://doi.org/10.1007/s10038-005-0237-9>.
- [153] M. Zaghlula, D.G. Glaze, G.M. Enns, L. Potocki, A.L. Schwabe, B. Suter, Current clinical evidence does not support a link between TBL1XR1 and Rett syndrome: description of one patient with Rett features and a novel mutation in TBL1XR1, and a review of TBL1XR1 phenotypes, *Am. J. Med. Genet. Part A* 176 (2018) 1683–1687, <https://doi.org/10.1002/ajmg.a.38689>.
- [154] C.A. Heinen, A. Jongejan, P.J. Watson, B. Redeker, A. Boelen, O. Boudzovitch-Surovtseva, F. Forzano, R. Hordijk, R. Kelley, A.H. Olney, M.E. Pierpont, G.B. Schaefer, F. Stewart, a S.P. van Trotsenburg, E. Fliers, J.W.R. Schwabe, R.C. Hennekam, A specific mutation in TBL1XR1 causes Pierpont syndrome, *J. Med. Genet.* 53 (2016) 330–337, <https://doi.org/10.1136/jmed-genet-2015-103233>.
- [155] G. Bracaglia, B. Conca, A. Bergo, L. Rusconi, Z. Zhou, M.E. Greenberg, N. Landsberger, S. Soddu, C. Kilstrup-Nielsen, Methyl-CpG-binding protein 2 is phosphorylated by homeodomain-interacting protein kinase 2 and contributes to apoptosis, *EMBO Rep.* 10 (2009) 1327–1333, <https://doi.org/10.1038/embor.2009.217>.
- [156] R.Y. Tweedie-Cullen, J.M. Reck, I.M. Mansuy, Comprehensive mapping of post-translational modifications on synaptic, nuclear, and histone proteins in the adult mouse brain, *J. Proteome Res.* 8 (2009) 4966–4982, <https://doi.org/10.1021/pr9003739>.
- [157] E.L. Huttlin, M.P. Jedrychowski, J.E. Elias, T. Goswami, R. Rad, S.A. Beausoleil, J. Villén, W. Haas, M.E. Sowa, S.P. Gygi, A tissue-specific atlas of mouse protein phosphorylation and expression, *Cell* 143 (2010) 1174–1189, <https://doi.org/10.1016/j.cell.2010.12.001>.
- [158] M.L. Gonzales, S. Adams, K.W. Dunaway, J.M. LaSalle, Phosphorylation of distinct sites in MeCP2 modifies cofactor associations and the dynamics of transcriptional regulation, *Mol. Cell. Biol.* 32 (2012) 2894–2903, <https://doi.org/10.1128/MCB.06728-11>.
- [159] D.H. Yasui, M.L. Gonzales, J.O. Aflatoon, F.K. Crary, D.J. Hu, B.J. Gavino, M.S. Golub, J.B. Vincent, N.C. Schanen, C.O. Olson, M. Rastegar, J.M. LaSalle, Mice with an isoform-ablating MeCP2 exon 1 mutation recapitulate the neurologic deficits of Rett syndrome, *Hum. Mol. Genet.* 23 (2014) 2447–2458, <https://doi.org/10.1093/hmg/ddt640>.
- [160] A. Bergo, M. Strollo, M. Gai, I. Barbiero, G. Stefanelli, S. Sertic, C.C. Gigli, F. Di Cunto, C. Kilstrup-Nielsen, N. Landsberger, Methyl-CpG binding protein 2 (MeCP2) localizes at the centrosome and is required for proper mitotic spindle organization, *J. Biol. Chem.* 290 (2015) 3223–3237, <https://doi.org/10.1074/jbc.M114.608125>.
- [161] J. Cheng, M. Huang, Y. Zhu, Y.J. Xin, Y.K. Zhao, J. Huang, J.X. Yu, W.H. Zhou, Z. Qiu, SUMOylation of MeCP2 is essential for transcriptional repression and hippocampal synapse development, *J. Neurochem.* 128 (2014) 798–806, <https://doi.org/10.1111/jnc.12523>.
- [162] D.J.C. Tai, Y.C. Liu, W.L. Hsu, Y.L. Ma, S.J. Cheng, S.Y. Liu, E.H.Y. Lee, MeCP2 SUMOylation rescues MeCP2-mutant-induced behavioural deficits in a mouse model of Rett syndrome, *Nat. Commun.* 7 (2016) 10552, <https://doi.org/10.1038/ncomms10552>.
- [163] L. Zocchi, P. Sassone-Corsi, SIRT1-mediated deacetylation of MeCP2 contributes to BDNF expression, *Epigenetics* 7 (2012) 695–700, <https://doi.org/10.4161/epi.20733>.
- [164] A. Guo, H. Gu, J. Zhou, D. Mulhern, Y. Wang, K.A. Lee, V. Yang, M. Aguiar, J. Kornhauser, X. Jia, J. Ren, S.A. Beausoleil, J.C. Silva, V. Vemulapalli, M.T. Bedford, M.J. Comb, Immunoaffinity enrichment and mass spectrometry analysis of protein methylation, *Technol. Innov. Resour.* 13 (2014) 372–387, <https://doi.org/10.1074/mcp.O113.027870>.
- [165] Z. Wang, N.D. Udeshi, M.O. Malley, J. Shabanowitz, D.F. Hunt, G.W. Hart, Enrichment and site mapping of O-linked N-acetylglucosamine by a combination of chemical/enzymatic tagging, photochemical cleavage, and electron transfer dissociation mass spectrometry, *Mol. Cell. Proteom.* 9 (2010) 153–160, <https://doi.org/10.1074/mcp.M900268-MCP200>.
- [166] A. Becker, P. Zhang, L. Allmann, D. Meilinger, B. Bertulat, D. Eck, M. Hofstaetter, G. Bartolomei, M. Hottiger, V. Schreiber, H. Leonhardt, M.C. Cardoso, Poly(ADP-ribosylation) of methyl CpG binding domain protein 2 regulates chromatin structure, *J. Biol. Chem.* 291 (2016) 4873–4881, <https://doi.org/10.1074/jbc.M115.698357>.
- [167] K.D. Kernohan, Y. Jiang, D.C. Tremblay, A.C. Bonvissuto, J.H. Eubanks, M.R.W. Mann, N.G. Bérubé, ATRX partners with cohesin and MeCP2 and contributes to developmental silencing of imprinted genes in the brain, *Dev. Cell* 18 (2010) 191–202, <https://doi.org/10.1016/j.dev-cell.2009.12.017>.
- [168] A. Guarda, F. Bolognese, I.M. Bonapace, G. Badaracco, Interaction between the inner nuclear membrane lamin B receptor and the heterochromatic methyl binding protein, MeCP2, *Exp. Cell Res.* 315 (2009) 1895–1903, <https://doi.org/10.1016/j.yexcr.2009.01.019>.
- [169] H. Kimura, K. Shiota, Methyl-CpG-binding protein, MeCP2, is a target molecule for maintenance DNA methyltransferase, Dnmt1, *J. Biol. Chem.* 278 (2003) 4806–4812, <https://doi.org/10.1074/jbc.M209923200>.
- [170] P.-F. Cartron, A. Nadaradjane, F. Lepape, L. Lalier, B. Gardie, F.M. Vallette, Identification of TET1 partners that control its DNA-demethylating function, *Genes Cancer* 4 (2013) 235–241, <https://doi.org/10.1177/1947601913489020>.
- [171] D.M. Murphy, P.G. Buckley, S. Das, K.M. Watters, K. Bryan, R.L. Stallings, Co-localization of the oncogenic transcription factor MYCN and the DNA methyl binding protein MeCP2 at genomic sites in neuroblastoma, *PLoS One* 6 (2011), e21436, <https://doi.org/10.1371/journal.pone.0021436>.
- [172] S.W. Long, J.Y. Ooi, P.M. Yau, P.L. Jones, A brain-derived MeCP2 complex supports a role for MeCP2 in RNA processing, *Biosci. Rep.* 31 (2011) 333–343, <https://doi.org/10.1042/BSR20100124>.
- [173] R. Li, Q. Dong, X. Yuan, X. Zeng, Y. Gao, C. Chiao, H. Li, X. Zhao, S. Keles, Z. Wang, Q. Chang, Misregulation of alternative splicing in a mouse model of Rett syndrome, *PLoS Genet.* 12 (2016), e1006129, <https://doi.org/10.1371/journal.pgen.1006129>.
- [174] S.G. Dastidar, F.H. Bardai, C. Ma, V. Price, V. Rawat, P. Verma, V. Narayanan, S.R. D'Mello, Isoform-specific toxicity of MeCP2 in postmitotic neurons: suppression of neurotoxicity by FoxG1, *J. Neurosci.* 32 (2012) 2846–2855, <https://doi.org/10.1523/JNEUROSCI.5841-11.2012>.

- [175] F. Mari, S. Azimonti, I. Bertani, F. Bolognese, E. Colombo, R. Caselli, E. Scala, I. Longo, S. Grosso, C. Pescucci, F. Ariani, G. Hayek, P. Balestri, A. Bergo, G. Badaracco, M. Zappella, V. Broccoli, A. Renieri, C. Kilstup-Nielsen, N. Landsberger, CDKL5 belongs to the same molecular pathway of MeCP2 and it is responsible for the early-onset seizure variant of Rett syndrome, *Hum. Mol. Genet.* 14 (2005) 1935–1946, <https://doi.org/10.1093/hmg/ddi198>.
- [176] A. Khoshnan, P.H. Patterson, Elevated IKK α accelerates the differentiation of human neuronal progenitor cells and induces MeCP2-dependent BDNF expression, *PLoS One* 7 (2012), e41794, <https://doi.org/10.1371/journal.pone.0041794>.
- [177] K.N. McFarland, M.N. Huizenga, S.B. Darnell, G.R. Sangrey, O. Berezovska, J.H.J. Cha, T.F. Outeiro, G. Sadri-Vakili, MeCP2: a novel huntingtin interactor, *Hum. Mol. Genet.* 23 (2014) 1036–1044, <https://doi.org/10.1093/hmg/ddt499>.
- [178] A. Becker, L. Allmann, M. Hofstätter, V. Casà, P. Weber, A. Lehmkühl, H.D. Herce, M.C. Cardoso, Direct homo- and hetero-interactions of MeCP2 and MBD2, *PLoS One* 8 (2013), e53730, <https://doi.org/10.1371/journal.pone.0053730>.
- [179] Q. Chang, G. Khare, V. Dani, S. Nelson, R. Jaenisch, The disease progression of Mecp2 mutant mice is affected by the level of BDNF expression, *Neuron* 49 (2006) 341–348, <https://doi.org/10.1016/j.neuron.2005.12.027>.
- [180] G.J. Pelka, C.M. Watson, T. Radziewicz, M. Hayward, H. Lahooti, J. Christodoulou, P.P.L. Tam, Mecp2 deficiency is associated with learning and cognitive deficits and altered gene activity in the hippocampal region of mice, *Brain* 129 (2006) 887–898, <https://doi.org/10.1093/brain/awl022>.
- [181] L.L.L. Orefice, A.L.L. Zimmerman, A.M.M. Chirila, S.J.J. Sleboda, J.P.P. Head, D.D.D. Ginty, Peripheral mechanosensory neuron dysfunction underlies tactile and behavioral deficits in mouse models of ASDs, *Cell* 166 (2016) 299–313, <https://doi.org/10.1016/j.cell.2016.05.033>.
- [182] A. Lawson-Yuen, D. Liu, L. Han, Z.I. Jiang, G.E. Tsai, A.C. Basu, J. Picker, J. Feng, J.T. Coyle, Ube3a mRNA and protein expression are not decreased in Mecp2R168X mutant mice, *Brain Res.* 1180 (2007) 1–6, <https://doi.org/10.1016/j.brainres.2007.08.039>.
- [183] L.R. Schaevitz, N.B. Gómez, D.P. Zhen, J.E. Berger-Sweeney, MeCP2 R168X male and female mutant mice exhibit Rett-like behavioral deficits, *Genes Brain Behav.* 12 (2013) 732–740, <https://doi.org/10.1111/gbb.12070>.
- [184] E. Wegener, C. Brendel, A. Fischer, S. Hülsmann, J. Gärtner, P. Huppke, Characterization of the MeCP2R168X knockin mouse model for Rett syndrome, *PLoS One* 9 (2014), e115444, <https://doi.org/10.1371/journal.pone.0115444>.
- [185] M.R. Pitcher, J.A. Herrera, S.A. Buffington, M.Y. Kochukov, J.K. Merritt, A.R. Fisher, N.C. Schanen, M. Costa-Mattioli, J.L. Neul, Rett syndrome like phenotypes in the R255X Mecp2 mutant mouse are rescued by MECP2 transgene, *Hum. Mol. Genet.* 24 (2015) 2662–2672, <https://doi.org/10.1093/hmg/ddv030>.
- [186] C.S. Casas-Delucchi, A. Becker, J.J. Bolius, M. Cristina Cardoso, Targeted manipulation of heterochromatin rescues MeCP2 Rett mutants and re-establishes higher order chromatin organization, *Nucleic Acids Res.* 40 (2012), <https://doi.org/10.1093/nar/gks784>.
- [187] E. Ballestar, T.M. Yusufzai, A.P. Wolffe, Effects of Rett syndrome mutations of the Methyl-CpG binding domain of the transcriptional repressor MeCP2 on selectivity for association with methylated DNA, *Biochemistry* 39 (2000) 7100–7106, <https://doi.org/10.1021/bi0001271>.
- [188] T.M. Yusufzai, A.P. Wolffe, Functional consequences of Rett syndrome mutations on human MeCP2, *Nucleic Acids Res.* 28 (2000) 4172–4179, <https://doi.org/10.1093/nar/28.21.4172>.
- [189] A. Free, R.I.D. Wakefield, B.O. Smith, D.T.F. Drydenll, P.N. Barlow, A.P. Bird, DNA recognition by the methyl-CpG binding domain of MeCP2, *J. Biol. Chem.* 276 (2001) 3353–3360, <https://doi.org/10.1074/jbc.M007224200>.
- [190] S. Kudo, M. Tamura, Y. Nomura, M. Segawa, N. Fujita, M. Nakao, J. Dragich, C. Schanen, Functional analyses of MeCP2 mutations associated with Rett syndrome using transient expression systems, *Brain Dev.* 23 (2001) 165–173, [https://doi.org/10.1016/S0387-7604\(01\)00345-X](https://doi.org/10.1016/S0387-7604(01)00345-X).
- [191] T. Baubec, R. Ivánek, F. Lienert, D. Schübeler, Methylation-dependent and -independent genomic targeting principles of the MBD protein family, *Cell* 153 (2013) 480–492, <https://doi.org/10.1016/j.cell.2013.03.011>.
- [192] J.R. Sinnamon, S.Y. Kim, G.M. Corson, Z. Song, H. Nakai, J.P. Adelman, G. Mandel, Site-directed RNA repair of endogenous Mecp2 RNA in neurons, *Proc. Natl. Acad. Sci. U. S. A.* 114 (2017) E9395–E9402, <https://doi.org/10.1073/pnas.1715320114>.
- [193] I. D'Annessa, A. Gandaglia, E. Brivio, G. Stefanelli, A. Frasca, N. Landsberger, D. Di Marino, Tyr120Asp mutation alters domain flexibility and dynamics of MeCP2 DNA binding domain leading to impaired DNA interaction: atomistic characterization of a Rett syndrome causing mutation, *Biochim. Biophys. Acta Gen. Subj.* 1862 (2018) 1180–1189, <https://doi.org/10.1016/j.bbagen.2018.02.005>.
- [194] R.P. Ghosh, R.A. Horowitz-Scherer, T. Nikitina, L.M. Gierasch, C.L. Woodcock, Rett syndrome-causing mutations in human MeCP2 result in diverse structural changes that impact folding and DNA interactions, *J. Biol. Chem.* 283 (2008) 20523–20534, <https://doi.org/10.1074/jbc.M803021200>.
- [195] B.J.O. Roak, L. Vives, W. Fu, J.D. Egerton, I.B. Stanaway, G. Ian, G. Carvill, A. Kumar, C. Lee, K. Ankenman, J. Munson, J.B. Hiatt, E.H. Turner, R. Levy, D.R.O. Day, N. Krumm, P. Bradley, B.K. Martin, E. Borenstein, D. a Nickerson, C. Heather, D. Doherty, J.M. Akey, R. Bernier, E.E. Eichler, J. Shendure, Multiplex targeted sequencing identifies recurrently mutated genes in autism spectrum disorders, *Science* 338 (2012) 1619–1622, <https://doi.org/10.1126/science.1227764>.
- [196] B.J.O. Roak, L. Vives, S. Girirajan, E. Karakoc, N. Krumm, P. Coe, R. Levy, A. Ko, C. Lee, J.D. Smith, E.H. Turner, B. Ian, B. Vernot, M. Malig, C. Baker, B. Reilly, J.M. Akey, E. Borenstein, M.J. Rieder, D. a Nickerson, R. Bernier, E.E. Eichler, Sporadic autism exomes reveal a highly interconnected protein network of de novo mutations, *Nature* 485 (2012) 246–250, <https://doi.org/10.1038/nature10989>.